

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Drug Resistance Updates

journal homepage: www.elsevier.com/locate/drug

Cancer Immunotherapy Getting Brainy: Visualizing the Distinctive CNS Metastatic Niche to Illuminate Therapeutic Resistance



Mark Owyong^a, Niloufar Hosseini-Nassab^b, Gizem Efe^a, Alexander Honkala^b,
Renske J.E. van den Bijgaart^c, Vicki Plaks^{d,*}, Bryan Ronain Smith^{b,*}

^a Department of Anatomy, University of California, San Francisco, CA 94143-0452, USA

^b Department of Radiology, Stanford University, Stanford, CA 94306, USA

^c Department of Radiation Oncology, Radiotherapy and Oncoimmunology Laboratory, Radboudumc, Geert Grooteplein Zuid 32, 6525, GA, Nijmegen, The Netherlands

^d Department of Orofacial Sciences, University of California, San Francisco, CA 94143, USA

A B S T R A C T

The advent of cancer immunotherapy (CIT) and its success in treating primary and metastatic cancer may offer substantially improved outcomes for patients. Despite recent advancements, many malignancies remain resistant to CIT, among which are brain metastases, a particularly virulent disease with no apparent cure. The immunologically unique niche of the brain has prompted compelling new questions in immuno-oncology such as the effects of tissue-specific differences in immune response, heterogeneity between primary tumors and distant metastases, and the role of spatiotemporal dynamics in shaping an effective anti-tumor immune response. Current methods to examine the immunobiology of metastases in the brain are constrained by tissue processing methods that limit spatial data collection, omit dynamic information, and cannot recapitulate the heterogeneity of the tumor microenvironment. In the current review, we describe how high-resolution, live imaging tools, particularly intravital microscopy (IVM), are instrumental in answering these questions. IVM of pre-clinical cancer models enables short- and long-term observations of critical immunobiology and metastatic growth phenomena to potentially generate revolutionary insights into the spatiotemporal dynamics of brain metastasis, interactions of CIT with immune elements therein, and influence of chemo- and radiotherapy. We describe the utility of IVM to study brain metastasis in mice by tracking the migration and growth of fluorescently-labeled cells, including cancer cells and immune subsets, while monitoring the physical environment within optical windows using imaging dyes and other signal generation mechanisms to illuminate angiogenesis, hypoxia, and/or CIT drug expression within the metastatic niche. Our review summarizes the current knowledge regarding brain metastases and the immune milieu, presents the current status of CIT and its prospects in targeting brain metastases to circumvent therapeutic resistance, and proposes avenues to utilize IVM to study CIT drug delivery and therapeutic efficacy in preclinical models that will ultimately facilitate novel drug discovery and innovative combination therapies.

I. Targeting brain metastasis with cancer immunotherapy

Brain Metastasis and their Unique Microenvironment

Metastasis, the spread of cancer cells from the primary tumor to surrounding tissues and distant organs, is the leading cause of cancer morbidity and mortality (Guan, 2015). Of particular interest are brain metastases, the treatment of which is a critical unmet need in order to successfully combat cancer. It has been estimated that brain metastasis occurs in up to 30% of patients across various solid cancers (Caffo et al.,

2013; Kyritsis et al., 2012). The most common source of brain metastases stems from non-small cell lung cancer (NSCLC), followed by breast, melanoma, renal, and colorectal cancers (Caffo et al., 2013; Kyritsis et al., 2012; Franceschini et al., 2016; Johanns et al., 2016).

The composition of the brain's microenvironment, specifically the immune milieu, is distinct from other tissues since it is immune privileged, which is attributable to the presence of a blood brain barrier (BBB) and the lack of conventional lymphatic drainage due to the absence of lymphatic fluid surrounding the brain (Louveau et al., 2015). Traditionally, lymphatic drainage enables circulation of maturing

* Corresponding authors.

E-mail addresses: vicki.plaks@ucsf.edu (V. Plaks), brsmith@stanford.edu (B.R. Smith).

¹ These authors contributed equally

antigen-presenting cells (APCs) to defined lymphatic structures where adaptive immune responses are preferentially mediated. The brain's unconventional lymphatic drainage presents an unclear anatomical route by which APCs traffic from the central nervous system (CNS) parenchyma to delineate the role of antigen presentation in neuro-inflammatory diseases. Within the brain, lymphatic fluid drains to the cervical lymph nodes through the subarachnoid space and ventricles by means of cerebrospinal fluid (CSF). In addition, the extracellular space of the brain and spinal cord parenchyma undergo lymphatic drainage by means of interstitial fluid (ISF) (Engelhardt et al., 2016). Another unique feature within the brain is the neurovascular unit (NVU), which consists of the BBB, endothelial cells and surrounding pericytes, astrocytes, neurons, and extracellular matrix (ECM). Heterogeneous NVU function and inflammation within the brain metastatic site often leads to inconsistent delivery of therapeutics and contrast imaging agents/modalities, a factor that must be considered in immuno-oncology (Puhalla et al., 2015).

The immune landscape of the brain during chronic inflammation is predominantly thought to consist of microglia, astrocytes, myeloid-derived suppressor cells (MDSCs), and tumor-associated macrophages (TAMs). Microglia, which are considered the primary APC in the brain microenvironment, serve as tissue-resident macrophages and may adapt to become perivascular macrophages that aid in tumor proliferation, invasion, and angiogenesis to create a more favorable tumor microenvironment (TME) (Hambardzumyan et al., 2016; Herz et al., 2017). It has been shown that microglia have the potential to differentiate into the M1-like or M2-like macrophages to induce a pro-inflammatory (TNF α /IFN γ response) or pro-tumoral (IL-4/TGF β response) phenotype, respectively (Lorger, 2012). Astrocytes, the most abundant glial cell type in the brain microenvironment, induce a pro-inflammatory response through the secretion of TNF α along with other cytokines (Constantinescu et al., 2005). Exhibiting multiple roles, astrocytes within the brain TME reduce survival of newly arriving metastatic cells, while promoting the growth of established brain metastases, highlighting one aspect of the unique tumor-stroma interaction of the brain (Quail and Joyce, 2017). Although these immune subtypes play various roles in the establishment of brain metastases, the infiltration of lymphocytes within the brain microenvironment is important in immuno-oncology since most approved cancer immunotherapy (CIT) drugs focus on enhancing T cell anti-tumor immunity. Despite the fact that the exact nature of the heterogeneous immune cell presence and mechanism of infiltration within the TME has thus far been elusive, partly due to the lack of imaging methods to visualize penetration across the BBB, preclinical data show that depletion of CD4 and CD8 T cells results in increased development of brain metastases and that regulatory T cells (Tregs), a subset of CD4 T cells that suppress anti-tumor immune responses, restrict the expansion and differentiation of T effector cells (Shevach, 2011; Lu et al., 2003). Understanding the dynamic interaction between the unique immune milieu and metastatic tumor cells within the brain TME will underpin which immune subsets are critical for immuno-oncology as therapeutic targets.

Current Diagnosis, Treatment, and Monitoring Options for Brain Metastases: Efficacy, Side Effects, and Challenges

The gold standard for diagnosis and monitoring of CNS metastases in the clinic is contrast-enhanced, high-resolution magnetic resonance imaging (MRI) (Puhalla et al., 2015). However, it is challenging to assess tumor response to a therapy and, more importantly, to differentiate between treatment-related changes or disease recurrence. This is due to the unique anatomy of the brain's vasculature, including the micro-environment of the NVU. Current treatment options for brain metastases include whole brain radiation therapy (WBRT), surgical resection, and stereotactic radiosurgery (SRS), all of which are altered depending on the size, number, histology, symptoms and location of metastatic lesions presented at the time of diagnosis (Lin and DeAngelis, 2015; Hatiboglu et al., 2013). WBRT remains the standard-of-care for treatment of metastatic CNS lesions, whereas surgical resection and SRS are considered effective for patients with manageable lesions (Johanns et al., 2016). Typically, WBRT and surgical resection are used for multiple or large lesions, while high-resolution MRI is used to assist with SRS for smaller, emerging lesions, and tumor treating fields (TTFields) are an emerging option for brain metastasis treatment (Staudt et al., 2010; Cohen and Kluger, 2016; Hottinger et al., 2016). However, each of these treatment options for brain metastases displays key limitations and consequences. WBRT can result in cognitive decline and SRS has been shown to be associated with radiation necrosis (RN), cerebral edema, and delayed tumor hemorrhage (Cohen and Kluger, 2016). Focal therapies, such as SRS and surgery, often have limited efficacy because of distant cerebral relapse and lack of treatment of microscopic tumor foci that remain invisible with current imaging technologies (Cohen and Kluger, 2016). TTFields, which disrupt cell division through the physical interactions of oscillating electric fields with key molecules during mitosis, have been applied only to glioblastoma multiforme (GBM) treatment thus far. They hold promise in brain metastasis to specifically address non-resectable regions of the brain and potentially affect stray cells leftover after resection or other interventions in brain metastasis. TTFields in combination with chemotherapy has shown promising early data in a phase 3 clinical trial of GBM in which prolonged progression-free survival and overall survival (OS) were noted in patients with recurrent disease (Hottinger et al., 2016). Notably, the non-invasive therapeutic potential of TTFields has been demonstrated by immense improvement of the OS of GBM, reducing tumor growth by up to 50% in preclinical studies alongside early clinical data in recurrent GBM demonstrating an increase in progression free survival after 6 months of treatment by 50% (Mittal et al., 2017). Despite the success of TTFields, they have resulted in seizures and nervous system disorders, such as anxiety and insomnia and although TTFields currently present the best quality of life for brain cancer patients, more evidence is needed to validate that other aspects of life quality do not suffer from the continual electric field stimulation required to achieve TTFields' cytotoxic effects (Hottinger et al., 2016).

Primary challenges when studying brain metastases include (Table 1): first, the lack of imageable preclinical models that accurately

Table 1
Features of Brain Metastasis Treatment (Johanns et al., 2016; Puhalla et al., 2015; Lin and DeAngelis, 2015; Hatiboglu et al., 2013; Staudt et al., 2010; Cohen and Kluger, 2016).

Current Treatment Modalities	Modality Side Effects	Associated Therapy Challenges
Whole-body radiotherapy (WBRT)	Radiation toxicity	Differentiation of primary and secondary tumor responses to treatment
Stereotactic radiosurgery (SRS)	Neurocognitive decline	Definition of tumor borders and micrometastases
Brachytherapy	Brain tissue damage	Heterogeneous inflammatory response of neurovascular unit
Surgical resection	Surgical trauma	Detection of minimal residual disease margins
Monitoring disease progression or treatment response via high-resolution CT or MRI	Repeated exposure to CT radiotoxicity and/or MRI contrast agent toxicity	Detection of quiescent micrometastases, differentiation of necrotic vs. inflamed cerebral edema, patient compliance

recapitulate CNS metastases for both mechanistic and therapeutic studies limits our knowledge of the immune contexture and the ability to accurately study pharmacodynamic parameters. A second challenge involves image standardization of brain metastases to assess response, as well as inclusion of novel imaging models that recapitulate early to late events of brain metastasis (Puhalla et al., 2015). Developing intravital microscopy (IVM) methods to effectively image the cellular and molecular determinants of standard-of-care and newly approved CIT drugs could facilitate the development of relevant preclinical models for data-driven combination therapy as well as provide insight into the design of more inclusive clinical trials. IVM offers the spatiotemporal resolution and multi-plexing potential to delineate the dynamic interactions of drugs, immune effectors, and tumor cells, which will lead to improved dosing and delivery of CIT drugs in monotherapy and provide a more rational foundation for testing combination therapies.

Would CIT be effective against brain metastases across various primary cancers?

Emerging systemic immune-modulating treatments have shown prolonged survival for patients with aggressive extra-cerebral disease, specifically brain metastasis from NSCLC and malignant melanoma (Owen and Souhami, 2014; Di Giacomo and Margolin, 2015). Although it had previously been thought that the brain TME is immune privileged and impervious to lymphocyte infiltration, increasing evidence has shown that tumor infiltrating lymphocytes (TILs) do traffic into the brain, where their presence is associated with increased overall survival and better response to CIT. Indeed T cells can cross the BBB, supporting the use of CIT in treating brain metastasis (Prins et al., 2006). Moreover, microglia, the resident macrophages of the brain, express PD-1 and activation of either PD-1, its ligand PD-L1, or CTLA-4 result in suppression of the anti-tumor T cell response (Berghoff et al., 2015; Pukrop et al., 2010). Of specific interest are anti-cancer TILs: CD3+, CD8+, and memory T cells, which have been shown to accumulate around the brain parenchyma along with immune-suppressive lymphocytes such as Tregs and PD-1-positive TILs (Berghoff and Preusser, 2015). However, to date there are no published results from clinical studies examining CIT pharmacokinetics or pharmacodynamics into brain lesions, primarily due to challenges in recruiting patients into trials requiring brain biopsies, particularly from patients responding well to therapies (Cohen and Kluger, 2016).

One main challenge currently being addressed by CIT is the study of whether CIT antibodies transit across the BBB despite their large size (Di Giacomo and Margolin, 2015; Chacko, 2013). Limited clinical evidence using positron emission tomography scans suggests that monoclonal antibodies are indeed able to cross the BBB, which in turn suggests that CIT drugs may be an effective means by which to target infiltrating TILs and microglial immuno-suppressive cell types (Ahluwalia and Winkler, 2015). Adsorptive transcytosis has also been evaluated as a potential vector for delivering drugs. This mechanism facilitates the transport of peptides through the interaction of a ligand with moieties expressed at the luminal surface of cerebral endothelial cells (Wilson et al., 2010; Hervé et al., 2008). However, this nonspecific transport process also occurs in other organs, complicating determination of effective drug doses in the brain, while reducing off-target drug dissemination. CIT drugs for brain metastasis have shown the most clinical promise in patients with melanoma and NSCLC; in particular, antibodies to CTLA-4 and PD-1 cross the BBB and evoke a partial response or in some cases stable disease (Margolin et al., 2012; NCT02085070; Goldberg et al., 2016). An additional challenge in the field of CIT for brain metastasis is the management of neurological symptoms during or after treatment that potentially arises from perilesional edema, intralesional hemorrhage, necrosis in previously irradiated lesions, or tumor growth due to treatment failure stemming from the inability to target microscopic tumor foci not evident from imaging. Post-immunotherapy side-effects have been addressed through the use

of surgery, radiation, steroids, or VEGF inhibitors to control edema, hemorrhage, and/or necrosis (Cohen and Kluger, 2016)

Finally, multidrug resistance (MDR) to anticancer drugs continues to hinder curative therapy of various human malignancies (Gottesman et al., 2016; Gonen and Assaraf, 2012; Zhitomirsky and Assaraf, 2016; Wijdeven et al., 2016; Bar-Zeev et al., 2017). MDR may arise from a variety of molecular mechanisms, such as new mutations in key target genes, dysregulation of normal apoptotic controls, or upregulation of chemoresistance due to multidrug efflux pumps of the ATP-binding cassette superfamily such as P-glycoprotein (P-gp), which expel a wide spectrum of cytotoxic agents (Li et al., 2016a). Interestingly, although it has been shown that P-gp is highly expressed within the BBB of humans, patients with brain metastases from malignant melanoma or lung cancer, present with lower P-gp expression than normal brain tissue, suggesting that there are other mechanisms of MDR functioning in metastatic brain tumors (Regina et al., 2001). With rapid advances in the understanding of MDR and CIT mechanisms, devising novel pre-clinical modalities that could inform on relevant clinical approaches to overcome the frequent emergence of cancer MDR (e.g. by directly targeting resistant cells with CIT) is a major need toward the goal to overcome tumor resistance to therapy.

Current progress in targeting brain metastasis with CIT

Checkpoint inhibition, specifically CTLA-4, PD-1, and its ligand PD-L1, function through the activation of CD4 and CD8 effector T cells. Typically, CTLA-4 and PD-1 signaling inhibit T cell activation, and blockade of either modulatory signal can shift the immune system's balance towards activation of T cells, thereby promoting tumor destruction (Buchbinder and Desai, 2016). While CD28 acts as a strong positive stimulatory receptor to CD80 and CD86, two T cell receptors that initiate and maintain CD4+ T cell proliferation, CTLA-4 functions as a potent inhibitory molecule against CD80 and CD86 (Vasilevko et al., 2002). PD-1/PD-L1, expressed by T cells, B cells, natural killer T (NKT) cells, activated monocytes, and dendritic cells, can also bind to CD80, which potentially forms a connection between the CD28/CTLA-4 and PD-1/PD-L1 pathways (Agata et al., 1996; Keir et al., 2008; Butte et al., 2007). More specifically, CTLA-4 is associated with inhibition of Akt signaling and PD-1/PD-L1 is associated with PI3K suppression and Akt signaling (Naidoo et al., 2014). PD-L1 is also upregulated on many tumor cells (Iwai et al., 2002). Signaling of PD-1 exerts a pro-tumorigenic effect by inhibiting production of IFN γ , TNF α , and interleukin-2, all of which are established anti-tumor response hallmarks (Riella et al., 2012). Moreover, chimeric antigen receptor T cell (CAR-T) therapy is increasingly gaining traction. The FDA's recent approval of Novartis' CAR-T therapy for B cell acute lymphoblastic leukemia and Kite Pharma's CAR-T therapy for non-Hodgkin's lymphoma demonstrates its promise for hematological malignancies. The field of immuno-oncology is heavily investing in CAR-T studies for solid tumors. CAR-T CIT involves the isolation of the patient's own T cells and genetically engineering them with a tumor-specific CAR onto their surface. This allows these modified T cells to specifically localize to and eliminate tumor cells by interacting with tumor-associated antigens (TAA) expressed on the tumor cell's surface (Yu et al., 2017). Although the structure of a CAR is similar to a T cell receptor, a CAR recognizes TAA independently of presentation via major histocompatibility complex (MHC) molecules, and targets a heterogeneous repertoire of antigens present on the tumor cell surface (Chmielewski et al., 2014; Fesnak et al., 2016). Table 2 details the FDA-approved CITs for specific cancer types and their performance in treating patients with brain metastasis. CIT has made groundbreaking strides particularly in hematological tumors and more recently in solid malignancies.

Preclinical studies with immune-modulating antibodies on primary CNS tumors have shown promising efficacy. Specifically, anti-CTLA-4 antibody was well tolerated in mice with SMA-650 intracranial tumors; a concomitant elevation of CD4+ cells and decrease in Tregs helped

Table 2

FDA Approved Immunotherapeutics and Current Progress on Brain Metastasis (Margolin et al., 2012; Goldberg et al., 2016; Tazi et al., 2015; Silk et al., 2013; Knisely et al., 2012; Larkin et al., 2015; Tan et al., 2017).

Immunotherapeutic	Target	Indicated Disease States	Current Progress on Brain Metastasis
Ipilimumab	CTLA-4	Unresectable or metastatic melanoma	Phase II clinical trials shows 10-25% disease control in metastatic brain cancer
Nivolumab	PD-1	Locally advanced or metastatic urothelial carcinoma	Phase III clinical trial extended survival of melanoma brain metastases patients by 6.9 months, while combination Nivolumab and Ipilimumab extended survival by 11.5 months
Atezolizumab	PD-L1	Metastatic non-small cell lung cancer (NSCLC) and locally advanced or metastatic urothelial carcinoma	Phase III clinical trial is currently on-going and shows promising preliminary results in patients with NSCLC and brain metastasis
Avelumab	PD-L1	Metastatic Merkel cell carcinoma (MCC)	Patients with brain metastasis are excluded from clinical trials using Avelumab
Durvalumab	PD-L1	Locally advanced or metastatic urothelial carcinoma	Phase II clinical trial currently recruiting participants with brain metastases from epithelial-derived tumors
Pembrolizumab	PD-1	Microsatellite instability-high (MSI-h) or mismatch repair deficient (dMMR) unresectable or metastatic solid tumors	Phase II clinical trial of brain metastases in melanoma and NSCLC currently on-going with early responses reported in 20-30% of patients
Chimeric Antigen Receptor T Cell Therapy (CAR-T)	Antigen on targetted tumor cells	B cell acute lymphoblastic leukemia (ALL) and non-Hodgkin's lymphoma (NHL)	Phase I clinical trial for brain tumors currently under-going recruitment

CD4 and CD8 T cell infiltration within the brain microenvironment is not well described, but preclinical data on brain metastases patients show promise in efficacy of checkpoint inhibition.

increase survival in these animals (Cohen and Kluger, 2016; Fecci et al., 2007). In other animal models, the combination of PD-1 and CTLA-4 inhibitors led to improved survival as compared to either single agent (Cohen and Kluger, 2016; Reardon et al., 2016). These studies indicate that BBB drug penetration might occur in both primary and metastatic CNS tumors, and clinical trials were undertaken to expand the studies on human subjects (Cohen and Kluger, 2016).

One of the first clinical studies evaluating the effect of CIT on brain metastases from melanoma was the phase II trial with ipilimumab (anti-CTLA-4), in which 12 of 115 patients randomized in the parent trial had stable brain metastases at baseline and were evaluated for efficacy. In the study, 2 of the 12 patients partially responded and 3 had stable disease, which were both alive at the last follow-up with a median patient survival of 14 months (range: 2.7-56.4+) (Weber et al., 2011; Weber et al., 2009). A different phase II trial with 72 melanoma patients suffering from brain metastases showed that ipilimumab prolonged OS and the OS was particularly significant in asymptomatic brain metastasis patients (Heller et al., 2011). Based on these promising results, a phase II trial of combination ipilimumab and fotemustine (NIBIT-M1) treatment on 20 asymptomatic patients with brain metastases was initiated. 25% of the patients had stable disease or partial response and another 25% had complete responses, indicating that antibody-based CIT may cross the BBB and in many patients drive durable control of brain metastasis (Di Giacomo and Margolin, 2015; Di Giacomo et al., 2015; Di Giacomo et al., 2012).

The potential for combining CIT following neoadjuvant radiotherapy to overcome therapeutic resistance in the treatment of brain metastasis

The field of CIT is actively shifting from monotherapy to combination treatment. However, whether this entails combining various CIT modalities to increase efficacy or to combine radio- or chemo-therapy with one or multiple CITs is currently debatable due to a lack of understanding of the changing immune milieu during and following therapy. However, it has recently been shown that the efficacy of immune checkpoint inhibitors can be enhanced by combining multiple inhibitors or via combination with chemotherapy or radiation therapy (RT) (Sindoni et al., 2017; Ott et al., 2017; Berghoff et al., 2016). This success can be attributed to a diverse T cell receptor repertoire of TILs following RT and CIT, alleviating the exhausted T cell phenotype, increasing the CD8/Treg ratios for effector function (Koyama et al., 2016; Curran et al., 2010; Pitt et al., 2017). In tumors resistant to RT and CIT,

a distinct increase in PD-L1 blunts the prevalence of CD8 TILs and interferes with T cell function (Twyman-Saint Victor et al., 2015). However, the absence of an effective innate immune system, including Toll-like receptor signaling (which is involved in antigen presentation), impairs the efficacy of radiotherapy-mediated control of tumorigenesis. Since radiation enhances the up-regulation of MHC class I in a dose-dependent manner to present peptide fragments on the cell surface to cytotoxic T cells, RT can trigger a systemic or local immune response. This response can elevate the expression of tumor-related antigens above a threshold level required for activation of circulating tumor lymphocytes, ultimately resulting in T cells potentially recognizing and attacking distant tumors (Reits et al., 2006). Irradiated cells also release a significant amount of high-mobility group box 1 protein, a potent pro-inflammatory mediator that activates dendritic cells and stimulates an immune response following radiation (Franceschini et al., 2016). Thus, both the innate and adaptive immune cells in the TME contribute to tumor cell death in the irradiated field.

A number of mechanisms has been elucidated that help explain the success of combination checkpoint inhibitors with radiation in a neoadjuvant setting. For instance, radiation upregulates inflammatory cytokines (e.g. TNF α , IFN γ , and CXCL16), which promote tumor detection and facilitate T cell infiltration (Cohen and Kluger, 2016). In some cases, CIT and RT can have a combined synergistic effect that leads to a systemic abscopal effect, or distant bystander effect, which refers to localized immune stimulation inciting a systemic immune response that results in tumor shrinkage at both treated and untreated sites. The effect thereby supports the use of radiation combined with immune-modulating agents. Studies on an orthotopic glioma mouse model showed that focal radiation therapy and CIT using anti-CTLA-4 improved survival via a CD4-dependent mechanism and generated antigen-specific memory that are likely valuable for long-term surveillance. Interestingly, although increases were seen in both CD8 and CD4 TILs, depletion of CD8 T cells had no effect on treatment outcomes while depletion of CD4 T cells abrogated the antitumor effect of RT with CIT (Belcaid et al., 2014). Another study using an intracranial implant of GL261, a mouse glioma cell line, showed a ~50% increase in survival when using RT with PD-1 where an immunogenic response including elevated CD8/IFN γ /TNF α was observed (Zeng et al., 2013). Collectively, recent work reveals that local RT can increase tumor-specific cytotoxic T-lymphocytes, induce the presentation of previously occult cancer antigens to T cells, and increase the permeability of the BBB to CIT agents.

What are the open questions in the field that could be addressed by IVM?

The major open questions involved in developing better therapies to treat and monitor brain metastases focus on understanding: 1) the efficiency of drug delivery through the BBB, 2) the mechanism of action of CIT within the brain, and 3) interactions with brain-specific stromal and immune cells, such as microglia or astrocytes. The primary limitation in imaging the brain is a lack of spatiotemporal resolution, which is compounded by: 1) heterogeneous NVU function and inflammation within the tumor, resulting in inconsistent imaging (Puhalla et al., 2015), 2) imaging modalities' inability to differentiate the enlargement of lesions due to inflammation, necrosis, or tumor growth (Cohen and Kluger, 2016), and 3) in both RT alone and in combination with CIT, it is difficult to differentiate RN from tumor progression by current imaging techniques (Verma et al., 2013; Colaco et al., 2016).

In this review we address how IVM can provide greatly improved imaging-based methodologies for diagnosis and monitoring of response to CIT in preclinical models. We describe how IVM on preclinical models can facilitate the development of new clinical imaging techniques that are pertinent for distinguishing the etiology of CNS lesions and provide insights into CIT's mechanism of action, efficiency, and mechanism of therapeutic delivery across the BBB to identify targets to improve therapeutic strategies.

II. Examining how intravital microscopy is used to study immune cells in brain metastasis pertinent to CIT drug delivery and action

IVM Techniques and Their Advantages

Methods including histological examination, biochemical assays, primary cell culture, and *ex vivo* microscopic approaches have been used for many years to study tumor growth and progression (Lee et al., 2013; Stoletov et al., 2013; Lee et al., 2006; De Yébenes et al., 1982; Degenhardt et al., 2006). While valuable, each of the above methods is limited to terminal endpoints and *ex vivo* data, and therefore cannot fully recapitulate the dynamics or complexity of cellular and subcellular interactions involved in tumor progression and metastasis at high resolutions. Such biases may lead to overlooking or even misinterpreting phenomena critical to the development of better cancer treatments

Table 3

Intravital Microscopy Techniques (Berning et al., 2012; Andresen et al., 2012; Oheim et al., 2006; Kim et al., 2000; Campagnola et al., 2002; Yun et al., 2003; Pope et al., 2014; Lin et al., 2013; Oron et al., 2004).

IVM Imaging Technique	Description	Penetration Depth	Resolution	
			Spatial	Temporal
Optical Frequency Domain Imaging (OFDI)	Advanced optical coherence tomography (OCT) tech that can rapidly examine tissue structure dynamics	3.8mm	~5 μm	~msec
Multi-Photon Fluorescent Microscopy (MPM)	Multiple photon fluorescence excitation enables higher penetration depths and novel imaging modes	300-600μm	~200nm	~msec
→ Second-Harmonic Generation (SHG)	Non-linear photonic imaging, label-free and discriminates highly ordered subcellular structures, e.g. Collagen	~500μm	~0.5 μm	~msec
→ Third-Harmonic Generation (THG)	Non-linear photonic imaging that is label-free and discriminates disordered structures, e.g. Lipid bodies	~300μm	~0.5 μm	~msec
Coherent Anti-Stokes Raman Scattering (CARS)	High-contrast, stimulated Raman-based spectroscopic imaging method that is label-free	200μm	~200 nm	~10 msec
1-Photon Fluorescent Microscopy (1P)	Typically using a confocal approach (pinhole) for fluorescence imaging	80-150μm	~200 nm	~msec
Fluorescent Lifetime Imaging Microscopy (FLIM)	Detection of molecular interactions based on differences in fluorescence decay rates	N/A	~nm	~500 fs




Fig. 1. *In-vivo* fluorescent multiphoton imaging of brain metastatic melanoma cells (red) interacting with brain vasculature (Tie2-GFP (green fluorescent protein) endothelial cells, green), creating a metastatic niche. The image was obtained from a depth of 200–210 μm. White arrows included in the cartoon overlay depict potential immune-tumor interactions that can be visualized using IVM, such as immune cell extravasation across the BBB, vascular permeability, and cell type distributions at endothelial and tumor borders. Image adapted from (Kienast et al., 2010).

(Abdulreda et al., 2013; Nobis et al., 2013). We summarize the current IVM imaging techniques in Table 3 to provide a more thorough understanding of the dynamic and high-resolution capabilities of IVM.

Advances in IVM offer high-resolution measurement of cell fate and behavior in living animal models, enabling long-term, dynamic studies of tumor growth and treatment effects. These advances include combination single- and multiphoton microscopy with advanced genetic models, improved surgical techniques, and fluorescent tracking of cell

fate for weeks to many months. Notably, compared with MRI and nuclear imaging modalities, as an optical modality IVM does not penetrate tissue deeply; moreover, the penetration depth of IVM is dependent upon the excitation and emission wavelengths as well as tissue type and size (Theer et al., 2003; Kobat et al., 2009; Helmchen and Denk, 2005). Yet the advantages of IVM help compensate for its deficiencies as depicted in Fig. 1. Many groups are further pushing the boundaries of what is possible in IVM, for example, developments that enable *in vivo* super-resolution—better than diffraction-limited spatial resolutions (Berning et al., 2012). Continued refinement of these techniques is revealing remarkable and previously unknown phenomena underlying the molecular, cellular, and functional biology of tumor growth, regenerative medicine, and immune cell trafficking (Fukumura et al., 2010). The ability to understand the interactions between tumor and immune cells within the tumor/metastatic microenvironments using preclinical models is key to effectively target and treat cancer with immunotherapies.

IVM of Tumors

Two different IVM imaging approaches are primarily used—chronic and acute (single-session) imaging—which tends to dictate the surgical techniques employed to access target tumor sites. The approach is selected based on the question to be answered (Fukumura et al., 2010; Hak et al., 2010). The surgical manipulation required to expose the area of interest for IVM can lead to inflammatory or other confounding effects. Thus, the architecture of the tissue of interest is a key constraint for the design of successful IVM experiments, in which minimally-perturbed tissue provides the most biologically-relevant results (Pittet and Weissleder, 2016). Acute preparations are not only limited in the duration and frequency of the observation, which may include irrevocable exteriorization of anatomy or surgery (Fukumura et al., 2010; Egeblad et al., 2008; Monsky et al., 2002; Wyckoff et al., 2007), but the preparation procedures might also impact the physiological parameters.

Chronic IVM often employs a clear, biocompatible window that is surgically implanted over the target site to enable long-term and repetitive microscopic access that may include the ability to stabilize motion (Smith et al., 2013; Smith et al., 2014). This method has been used at a number of sites, including the cranium. Cranial windows, an optically transparent glass placed over the brain, are appropriate for studying neural activity, brain blood flow, or brain metastasis, and may endure for several months up to a year as compared to, for example, dorsal and abdominal windows, which typically last 3–6 weeks (Alieva et al., 2014; Smith and Gambhir, 2017). Use of fluorescent probes, transgenic mouse models, and repeated imaging with IVM enables long-term, high-spatiotemporal resolution (down to $\sim 0.1 \mu\text{m}$ and up to hundreds of frames per second) observation of tumor progression at the cellular to subcellular level. Prudent choice of which cells and molecules to label with exogenous probes or to engineer as transgenic reporters can enable significant insight into cell identity, state, movement, and interaction. For example, fluorescent labeling of adoptively transferred CAR-T cells means that cell migration, infiltration, proliferation, and activation may be monitored and quantified dynamically, potentially in combination with a transgenic reporter model. This approach permits direct examination of T cell-tumor interactions so that the influence of cancer proximity on each CAR-T parameter may be directly measured. Moreover, microendoscopic methods combined with IVM provide access to deeper tissues for chronic imaging, such as the hippocampus or striatum, which cannot be directly observed via conventional means (Barretto et al., 2011). Therefore, with IVM, a deep-lying glioma may be tracked at high-resolution over months to measure, for example, dynamic three-dimensional angiogenesis progression, changing microcirculatory velocities, infiltration, and proliferation, and health of fluorescently-labeled immune cells interacting with the tumor (Barretto et al., 2011; Brown et al., 2001; Wu et al., 1992; Breart et al., 2008; Boissonnas et al., 2007).

IVM to Investigate Tumor Immunobiology

Many features of the TME, elaborated upon below, are amenable to interrogation via IVM. Apart from malignant cells, the TME contains various components that play a major role in influencing the outcome of the malignancy. These can be broadly classified into three main groups: cells of haematopoietic origin (cells of the immune system), cells of mesenchymal origin (including fibroblasts, myofibroblasts, mesenchymal stem cells), and the non-cellular components of the ECM, which consists of complex interacting proteins, glycoproteins and proteoglycans (Pattabiraman and Weinberg, 2014; Whiteside, 2008). The TME bidirectionally and strongly influences the effects of CIT and other therapies, making it an important parameter to study in concert with therapeutic delivery.

1. Tumor Growth

Use of transgenic mice and stably-transfected fluorophore-expressing tumor cell lines enables long-term observation of the dynamic growth of tumors. By observing the cellular morphology and subcellular processes involved in tumor growth in various tissue sites under treatment, IVM offers rapid insights into not only the proliferation and apoptosis of tumor cells, angiogenesis, and trafficking of immune cells, but also the effects of treatment. For example, tumor-focused IVM allows visualization of the efficacy of new chemo- or immuno-therapeutics as well as their multifactorial mechanisms of action (Frieboes et al., 2013; Pons-Salort et al., 2012). Notably, tumors do not necessarily need to express fluorophores, as label-free OFDI or CARS may also be used to measure parameters such as the rate and extent of tumor growth through tissue derangement (Vakoc et al., 2009).

2. Extracellular Matrix (ECM)

Aggressive lesions actively degrade and subvert the surrounding ECM to facilitate tumor growth, tissue invasion, and metastasis. MPM and SHG are well-suited for the *in vivo* study of tumor-mediated derangement of the ECM. As displayed in Table 3, SHG microscopy is highly sensitive to changes in collagen fibril/fiber structure that occur during cancer and can provide important biological information about ECM alterations that accompany cancer progression and metastasis. Hence, SHG can be used to quantify collagen fiber formation, direction, and remodeling that may be affected by cell signaling or mechanical/biomechanical interaction between cells and collagen fibers (Keikhosravi et al., 2014). Moreover, rapidly growing tumor cells directly and indirectly, through recruited immune cells, degrade the ECM through the secretion of matrix metalloproteinases (Kessenbrock et al., 2010). Detachment of tumor cells from the surrounding tissue structure precedes invasion and metastasis, and the establishment of new metastases requires degradation of ECM to establish a hospitable growth niche (Deryugina and Quigley, 2006), making it a critical tool to study the formation of the brain metastatic niche. Thus, IVM-based observation of tumor-mediated changes in ECM and tumor and angiogenic development offers the potential for profound insights into the impact of therapeutics.

3. Tumor Microenvironment (TME)

Tumor growth is fueled by both intrinsic mutational and transcriptional drivers within the tumor itself as well as by extrinsic environmental factors that arise through the interaction of tumor, stromal, and immune cells (Joyce and Fearon, 2015). Collectively, these extrinsic factors comprise the TME. Rapid tumor proliferation results in the secretion of glycolytic metabolism byproducts, depletion of local normoxia, cytokine signaling, and accumulation of damaged cancer and stromal cells (Biswas, 2015; Rodriguez et al., 2017; Sukumar et al., 2017). These features drive localized

inflammation that, in turn, results in the infiltration of diverse innate and adaptive immune cell subpopulations. Infiltrating anti-cancer immune effectors such as CD8+ and CD4+ T cells, M1 macrophages, and NKT cells coordinate tumor lysis and adaptive immune memory to tumor antigens. However, the infiltration/maturation of immunosuppressive cells, such as MDSCs and Tregs, into the TME results in potent inhibition of anti-cancer immune effectors (Joyce and Fearon, 2015; Galon et al., 2006). Increased immunosuppressive cellular infiltration significantly correlates with rapid disease progression, treatment resistance, and greater metastatic dissemination in a number of tumor types, including breast and skin cancer (Sica and Bronte, 2007; Diaz-Montero et al., 2009). These phenomena are further complicated by their interactions with a) tissue-specific stromal factors, such as the astrocytes and microglia of NVU in brain metastases, b) ongoing angiogenesis, c) lysyl oxidase- and matrix metalloproteinase-mediated ECM alterations, and d) the profound effects of chemo-, radio-, and immuno-therapy upon the TME itself (Zitvogel et al., 2008; Bracci et al., 2014). Furthermore, growing tumors are not governed by the architecture of their surrounding tissues and may create dense desmoplastic networks that result in widely varied circulation, drug infiltration, and interstitial pressure. Not only does the nature of stromal activation and immune infiltration vary by primary tumor tissue site, but metastases also form their own distinct TME upon arrival in distant tissues, which likely differ from the primary TME in remarkable and as-yet-undefined manners (Massague and Obenauf, 2016; Peinado et al., 2017). The influences of brain-specific factors, such as the BBB, microglia, or reduced lymphatic drainage, on the formation and evolution of brain metastasis TMEs are currently unknown and the results of their enhanced understanding may represent compelling targets for therapy development.

Transgenic animal studies, in which tumors or immune cells have been selectively edited, have been critical to define the mechanisms and interactions underlying the relationship of the tumor to its TME in previous literature (Zhou et al., 2011; Casey et al., 2015; Walrath et al., 2010). Combination of transgenic animal models, fluorescently-tagged antibodies, gene fusion reporter cell lines, and the spatiotemporal capabilities of IVM will yield revolutionary insight into the underlying immunobiology of tumor-rejecting and tumor-promoting immune responses to brain metastases, the mechanism of CIT in the brain, and the impact of NVU-specific microglial or astrocytic activity on therapeutic response. TME physico-chemical properties may also be imaged via exogenously-administered contrast agents, e.g., molecular probes with optical properties that vary upon change in pO₂ or pH, which are well-suited for *in vivo* study of the dynamics of tumor-mediated hypoxia and the downstream metabolic reactions of tumor stroma and infiltrating immune cells (Dellian et al., 1996; Helmlinger et al., 1997). IVM of the immune component of the TME is detailed below.

4. Blood Supply

OFDI and MPM may be used to measure the growth of new endothelial venules and capillaries associated with tumor growth, making them well-suited to characterize the mechanisms by which metastases establish initial blood supplies. IVM has also been used to directly measure hemodynamics, vascular permeability, vessel pore cutoff size, leukocyte-endothelial cell interactions, lymphangiogenesis, and circulation of investigational drugs into the tumor site (Nobis et al., 2013; Zipfel et al., 2003; Brown et al., 2001; Hoshida et al., 2006; Maurin et al., 2011; Ricard and Debarbieux, 2014). For example, IVM has helped demonstrate that chemotherapeutic efficacy in primary brain tumors and metastasis is extremely limited due to poor transport of the fluorescently-labeled drug across the BBB and blood brain tumor barrier (Ryu et al., 2012; Giedt et al., 2013; Vishwanath et al., 2009). This suggests that IVM could similarly be used to iteratively and directly test newly designed drugs for improved transport kinetics.

5. Immune Cell Infiltration

Tumor progression is intimately shaped by several layers of feedback with the tumor-responsive immune system, mandating examination of the infiltrating and *in situ* proliferating immune system in the study of the mechanisms underlying tumor growth. *In vivo* discrimination between unlabeled immune and non-immune cells in tissues is complex, so the best-suited approaches for study of immune infiltration are exogenously-administered (adoptively-transferred and/or fluorescently-labeled) or endogenously-expressed (transgenic-fluorescent immune cells) reporters (Abdulreda et al., 2013; Torcellan et al., 2017; Osswald and Winkler, 2013; Moalli et al., 2015; Deguine et al., 2010; Chtanova et al., 2014). Strategic use of Cre recombination to generate lineage- and function-specific conditional fluorescent reporters enables the detailed tracking of cell fate and migration through living tissue by cell type, such as tracking the chemotaxis of labeled stem cells through developing lung tissue (Desai et al., 2014). IVM may be used to examine the infiltration of immune cells into the TME, their interactions with the growing tumor, interactions between immune cell types, and immune influence on stromal cells and migratory cancer cells. Extension of these capabilities of IVM together with advanced exogenously-administered and endogenously-expressed probes for the study of immune function in the TME will likely yield significant insight into the underlying immunobiology of tumor-rejecting and tumor-promoting immune responses and could help guide the development of new immunomodulatory drugs.

IVM in Brain Immuno-Oncology

IVM is well-adapted to study tumor immunobiology and has been used in preclinical models of GBM to measure the synergistic effects of anti-angiogenic and cytotoxic therapy (Jarzyna et al., 2010; Karagiannis et al., 2017). Many features of the brain TME including, but not limited to, immune cell infiltration and interaction with CIT drugs are depicted in Fig. 2. IVM techniques promise to surpass alternative terminal, static immune response characterization methods by applying the capabilities to chronically study immune dynamics in response to tumor growth and treatment. The nature and interplay of anti- and pro-tumor immune effectors are critical determinants of treatment success, so a more detailed understanding of their underlying mechanisms, and how those mechanisms interact with the unique microenvironment of brain metastases, is necessary. The traditional static methods may alter the behavior or expression of immune or cancer cells upon sample collection and processing. Complementing these methods, or potentially in their stead, IVM offers dynamic, long-term, and high-resolution imaging of cell-cell interactions, tissue structure, and treatment effects in a minimally-perturbed (e.g., after the post-window implantation/surgical rest period) living animal, where the influences of distal tissue sites upon, and cell interactions within, the measured site remain intact.

Specifically, IVM can facilitate exploration of the following:

1. Fundamental Tumor Immunobiology

Use of cranial windows enables direct, dynamic imaging of cell-cell interactions, cell proliferation, and cell migration in and around metastatic and primary brain tumors (Kienast et al., 2010). Fluorescent transgenic reporters and/or exogenously-administered, labeled cells may be applied to study several properties fundamental to tumor growth and immune response, such as T cell activation, the effects of hypoxia on myeloid cell development in the tumor margin, or the spatial organization and migration rates of stromal and myeloid cells involved in tumor-promoting angiogenesis (Murdoch et al., 2008).

2. Checkpoint Inhibitor Development

As discussed above, monoclonal antibody-mediated neutralization of immune checkpoints has shown clinical benefit in various solid




Fig. 2. The brain metastatic microenvironment is shaped by tumor growth, immune infiltration, and stromal cell interactions. Several features, described clockwise, can be assayed via IVM (Fukumura et al., 2010; Brown et al., 2001; Wu et al., 1992; Vakoc et al., 2009; Osswald and Winkler, 2013; Fukumura and Jain, 2007; Gordon et al., 2017; Kamoun et al., 2009; Ameer-Beg et al., 2002; Witte et al., 2011; Jain, 2002; Ohkubo et al., 1991; Melder et al., 1995; Ardeshtirpour et al., 2012; Provenzano et al., 2008; van Munster and Gadella, 2005; Le et al., 2009; Burke and Brown, 2014; Tserevelakis et al., 2014; Weigelin et al., 2016; Kurokawa et al., 2015; Choi et al., 2012; Vakoc et al., 2005; Dietzel et al., 2014; Gershanov et al., 2017; Sun et al., 2009; Uckermann et al., 2014; Evans and Xie, 2008; Ellenbroek and van Rheenen, 2014; Fridman et al., 2017).

- A) Lymphangiogenesis reshapes lymphatic drainage and immune infiltration and may be assayed via MPM or OFDI
- B) Cell migration through lymphatic vessels can be observed by fluorescently labeling the cell types of interest to image via MPM
- C) The organization and proliferation of immune cells in the TME shapes host response and tumor growth, these processes have been observed with 1P and MPM but may be amenable to further investigation via OFDI, FLIM, and THG
- D) Changes in the ECM dictate tumor growth and cell interactions and may be imaged via FLIM, SHG, or potentially even THG
- E) Tumor growth and migration can easily be tracked with fluorescent reporter tumor cell lines and any IVM imaging modality
- F) Immune and microglial infiltration from endothelial vessels, particularly the special architecture of the BBB, can be imaged via 1P or MPM and may be detectable via THG
- G) NVU-specific features, such as neurons, astrocytes, and pericytes, may be observed via fluorescent labeling suggested above
- H) Angiogenesis is a critical process for tumors to continue growing, where blood vessel change can be imaged via contrast dyes and fluorescent antibodies against endothelial markers
- I) The rate of blood flow and pO2 levels can be imaged directly via 1P, MPM, and may be amenable to imaging via FLIM, SHG, or THG
- J) Immature blood vessels frequently present with enhanced permeability, which may be imaged via fluorescent conjugates of dextran to separate active extra/intravasation from passive diffusion in immune or cancer cell trafficking
- K) The BBB is a unique feature of the brain and may be imaged via fluorescent reagents and 1P, MPM, or CARS and may be suitable for imaging via OFDI, FLIM, SHG, or THG
- L) High endothelial venules are mature, selectively permeable blood vessel structures that may regulate the trafficking of immune and cancer cells around tumors, tracking their distribution and function via 1P and MPM, or other imaging techniques, will reveal critical trafficking parameters of metastasis initiation, survival, and dissemination.

tumors, such as melanoma, colorectal, lung, and urothelial bladder cancers (Li et al., 2016b). However, the development of new checkpoint inhibiting antibodies is hampered by the heterogeneity of the immune response and sample processing-induced changes in immune function in most currently-employed preclinical models of cancer. Intravenous and intratumoral injection of fluorescently-labeled monoclonal antibodies for proteins of interest enables intravital imaging of: 1) therapeutic target and immune effector distribution, 2) target dynamics, 3) cell type expression of target antigens, 4) function of labeled antigens in cell-cell interactions, and 5) infiltration of antibodies, such as anti-PD-L1 into the TME (Mikucki et al., 2015).

3. Cell Therapy Development

Cancer-targeted, adoptive cell therapy, such as CAR-T, is most effective when transferred cells survive and proliferate in the TME (Jaspers and Brentjens, 2017; Newick et al., 2017). Adoptive cell therapies rely upon isolation, modification, and expansion of isolated patient-derived immune cells to increase their affinity for TAAs. The *ex vivo* modification of immune cells presents an opportunity to fluorescently label cells to track their fate and activity *in vivo*. IVM may also be used to track the migration paths and antitumor activity of individual immune cell subtypes.

4. Combination Therapy

Combination therapy is a rapidly advancing sub-field of clinical oncology as described in Section I. Accumulating evidence, including a clinical trial targeting GBM, shows promise in RT with CIT drugs targeting PD-1/PD-L1 such as pembrolizumab, nivolumab, atezolizumab, and

durvalumab lead to enduring responses (Weichselbaum et al., 2017). While combination therapy shows immense promise in NSCLC and melanoma, careful therapeutic design consideration must be applied to the brain, because it is a sensitive organ, in order to leave healthy tissue intact during therapy. Immune checkpoint blockers such as CTLA-4 and PD-1/PD-L1 enhance the antitumor response induced by radiation therapy (Twyman-Saint Victor et al., 2015; Zeng et al., 2013; Fang et al., 2017; Bang et al., 2017; Demaria et al., 2005; Yoshimoto et al., 2014; Pilon-Thomas et al., 2010; Koo and Kim, 2016); thus, the interplay between the biological effects of RT and the immune system may be a critical mechanism for inhibiting tumor growth and extending OS in brain metastasis patients. However, the ability to successfully design combination therapeutic strategies is limited due to a lack of understanding of the molecular and cellular mechanisms in radio-oncoimmunology (Wu et al., 2017). IVM's spatiotemporal resolution overcomes limitations that constrain other techniques, enabling understanding of immune milieu changes following local or systemic treatment. Since tissue-resident immune cells cannot easily be studied *ex vivo* due to their short lifespan and inadequate recapitulation of the microenvironment, dynamic microscopic imaging of the immune and tumor milieus can critically contribute to uncover effective therapeutic combinations to target brain metastases.

III. Perspective: Utilizing IVM of brain metastasis to advance drug discovery and facilitate effective targeting with approved CIT

Characterization of Brain Metastatic Tumor Cells

Circulating tumor cells often accumulate at vascular branch points, penetrate through vascular walls, and remain in close proximity to microvessels in the process of remote niche formation (Pittet and

Weissleder, 2011). Recently, the fates of metastasizing cancer cells were tracked in relation to blood vessels in living mouse brains using real-time MPM, a common form of IVM, which revealed the single steps of metastasis formation *in vivo* over minutes to months (Kienast et al., 2010). Exploiting this strategy to observe metastasis formation under treatment conditions has key implications for the design of improved therapies, specifically by examining immunosuppression patterns and immune cell interactions. Multiple steps of metastasis depend on myeloid cells, such as escorting of circulating tumor cells out of the vasculature and into their niche in the brain—this rare event can now be visualized, and interventions against it can be studied. The use of stably-transfected fluorescent tumor cell lines and IVM techniques such as OFDI, MPM, or CARS, can also allow visualization of the distinct characteristics of brain metastasis vs. primary brain tumors, most notably GBM, and their differential immune milieus. Through the use of fluorescent labels, IVM could be further extended to study the infiltration (migration tracking, ECM disruption, and angiogenesis) and survival (tumor proliferation, immune cell capture, and tissue invasion) of incipient brain metastases from other origin tissues, where illumination of origin tissue-specific differences may yield actionable insight into tailored clinical treatment. The almost limitless potential to characterize various poorly understood aspects of brain metastases, such as the high propensity of melanomas to form brain metastases compared to other primary cancer types or the rapid development of brain metastases in individuals with lung carcinoma, can provide novel insights and opportunities to effectively treat, and even possibly cure, previously untreatable metastatic malignancies.

Interactions and Changes in TME

Although current preclinical and preliminary clinical data suggest that CIT and RT synergize in the treatment of brain metastases, several remaining open questions must be addressed to guide future clinical practice. These questions include whether, and how, infiltration of immune cells depends upon extravasation through the BBB, where brain endothelium, pericytes, and microglia may all exert brain-specific effects on infiltrating immune cells—particularly under the influence of a malignancy. It is also not well-understood how RT affects the brain microenvironment and increases the efficacy of CIT; evidence suggests that RT induces inflammatory responses in tissue, to which non-tumor-suppressed immune cells respond with greater vigor—an effect yet to be shown directly (Tabatabaei et al., 2017). In-depth understanding of how and why tumor-immune cell interactions and the vasculature are altered is crucial for improved prognosis and treatment decisions in the clinical setting. Furthermore, there are several other poorly-understood phenomena that may underlie the behavior and/or success of metastases in the TME, including, but not limited to: a) the interactions of pericytes/podocytes with invasive tumor margins, b) interactions of microglia with infiltrating lymphocytes and innate immune cells, c) the role of lymphangiogenesis, and d) immune cell-immune cell interactions. By exploiting the multiplexing capabilities of intravital imaging (i.e., using multiple fluorescent channels in parallel or in series) and its spatiotemporal resolution, the changes in the TME and immune cell infiltration resulting from treatment with different combinations of RT and CIT can be tracked, quantified, and understood. For example, MPM could readily be used to track the localization and trafficking of fluorescently-labeled myeloid cells in a metastatic brain tumor-bearing mouse model to examine the movement and co-localization of myeloid effectors in relation to tumor cells. This system could be further extended with an inducible Cre recombinase model to knockout putative immuno-suppressive tumor factors or immune-activating stromal factors to confirm the role of such factors in metastatic growth. In another instance, transient antibody labeling of checkpoint sites could be used to assess changes in the CIT “druggable state” after treatment with chemo- or radio-therapy. Potentially more interesting is dynamic observation of CIT within the brain TME, including whether

immunotherapeutics interact more with the classical CD4/CD8 T cells or with myeloid cells (e.g., perivascular macrophages) within the brain by employing OFDI or MPM with a fluorescently tagged CIT drug. Identification of the particular cells interacting, their lengths of interaction, and the tissue sites of such interactions may pave the way toward the development of new blocking agents to inhibit specific cell-cell interactions or provide hints toward a promising combination therapy. This will help address current challenges in the field of CITs such as the efficiency of drug delivery and mechanism and efficacy of action and dependence on drug size.

Tracking Immune Cells

Cognitive function, especially memory, is severely affected by brain metastases and the complications of treatment. For example, WBRT causes damage within the hippocampus (Dye et al., 2015). IVM can be used to study these effects because of its high spatiotemporal resolution, allowing tracking of the changes in immune cell interactions in models of cognitive dysregulation. This will provide novel insights into changes within the brain's immune milieu. This technique can be further combined with optogenetic switches, which enables the activation or repression of transgenic loci with specific wavelengths of light to directly study the effects of WBRT or other treatment on normal or diseased neural activity and whether CIT affects these functions.

Additionally, MPM performed on fluorescently labeled CAR-T cells in the brain (see Section II) could help uncover the multi-dimensional processes by which CAR-T cells function in the brain. Similarly, MPM could allow study of mechanisms of CAR-T exhaustion in solid tumors. Direct examination of adoptively transferred, fluorescently-labeled CAR-T cell-tumor interactions will thus yield insight on the major immunosuppressive molecules with the brain TME.

Drug delivery and mechanism of action. Current challenges in drug development for the treatment of brain metastases include the following:

- Accurate study of drug penetrance across the BBB as well as drug uptake and effect by cell type
- Differences in the mechanism of action of CIT drugs in the specific context of the brain TME (e.g., compared with primary brain tumors or with other metastatic sites)
- Altered expression patterns of immune checkpoints in the brain
- Brain-specific patterns of response to radio-, chemo-, and immunotherapies

IVM should be used to explore BBB integrity and diffusion or transport of small molecules across the endothelial-pericyte barrier using fluorophore-tagged drugs or proxy molecules, such as fluorescent high molecular-weight dextrans or nanoparticles of varied sizes and shapes (Muradashvili et al., 2012; Koffie et al., 2011; Smith et al., 2012). Using super-resolution techniques, IVM might support, for example, the direct identification of drug entry mechanism, i.e., whether therapeutics enter via transcytosis, interendothelial gaps, or other processes in the brain vasculature. This knowledge is critical to therapeutic design considerations. Drug uptake as a function of cell type may be assessed through a combination of exploration of trans-BBB transport and diffusion with fluorescently-labeled cells and may be further extended to assess the drug's effect on cell proliferation, survival, hypoxia, localization, or functional activity. Data gathered from such studies may inform whether brain-specific differences in the action of immunomodulatory drugs exist and affect the target therapeutic window. As noted above, microglia may express PD-1 in inflammatory conditions, but the effects of that expression on the function of infiltrating lymphocytes, pericyte, neuron, and astrocyte components of the NVU remain unknown. Better elucidation of the effect of PD-1 inhibition thereof in the unique context of the NVU and brain TME will enable design of more incisive CIT therapies in brain metastasis.

Furthermore, transient labeling of proteins/receptors using fluorophore-bound antibodies may be employed to image the spatio-temporal distribution of checkpoints targeted by checkpoint inhibitor therapy in brain metastases before, during, and/or after treatment. Mapping checkpoint expression may then guide the choice of combination therapies, dose levels, and dose scheduling in subsequent studies to combat the spread of brain metastases. By employing the above methods alone, or in combination with fluorescent protein-expressing tumors, the long-term effects of chemo-, radio-, and immuno-therapy on brain metastasis growth patterns, survival, and spread can be integratively studied. The use of IVM in these studies provides a unique advantage through long-term dynamic imaging with high spatio-temporal resolution in which endogenous tissue architecture and behavior are maintained in far greater detail than is possible with terminal endpoints and assays that derange tumor or immune behavior.

Continued progress in application of CIT to brain metastases will demand improved study design, for which IVM techniques are well-suited. Previously unanswerable questions, such as the migration dynamics of PD-1-positive microglia or the influence of CTLA-4 inhibition upon T cells transiting the BBB, may begin to be addressed through the long-term studies enabled by IVM. Such studies will provide pivotal proof for CIT mechanism of action and therapeutic efficacy in the unique TME of brain metastases. Furthermore, correlative studies wedding dynamic IVM use with traditional study endpoints, such as flow cytometry and immunohistochemistry, offers enhanced understanding of the cell-cell interactions that underlie brain metastasis progression and effective treatment. Combination of IVM with emerging terminal endpoints, such as high-dimensional cytometry or immunohistochemistry capable of surveying dozens of antigens (Yao et al., 2014; Pyne et al., 2009; Qiu et al., 2011; Kennedy-Darling et al., 2016), is also a compelling line of investigation that may provide revolutionary—and clinically actionable—insights into the spatiotemporal dynamics of immune and brain metastasis cell interactions that lead to better therapies for metastatic brain cancer patients.

Acknowledgments

B.R.S. acknowledges support from a K99/R00 award (K99 CA160764), an AACR Translational Breast Cancer Grant, and the Ben and Catherine IVY Foundation Grant. V.P. acknowledges support from the Cancer League and the California Breast Cancer Research Program (23IB-0018).

References

- Abdulreda, M.H., Caicedo, A., Berggren, P.O., 2013. Transplantation into the anterior chamber of the eye for longitudinal, non-invasive *in vivo* imaging with single-cell resolution in real-time. *J. Vis. Exp.* 73, e50466.
- Agata, Y., Kawasaki, A., Nishimura, H., Ishida, Y., Tsubata, T., Yagita, H., et al., 1996. Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. *Int. Immunol.* 8 (5), 765–772.
- Ahluwalia, M.S., Winkler, F., 2015. Targeted and immunotherapeutic approaches in brain metastases. *Am. Soc. Clin. Oncol. Educ. Book* 67–74.
- Alieva, M., Ritsma, L., Giedt, R.J., Weissleder, R., van Rheenen, J., 2014. Imaging windows for long-term intravital imaging: General overview and technical insights. *Intravital* 3 (2), e29917.
- Ameer-Beg, S., Barber, P., Hodgkiss, R., Locke, R., Newman, R., Tozer, G., et al., 2002. Application of multiphoton steady-state and lifetime imaging to mapping of tumour vascular architecture *in vivo*. pp. 85–95.
- Andresen, V., Pollok, K., Rinnenthal, J.L., Oehme, L., Gunther, R., Spiecker, H., et al., 2012. High-resolution intravital microscopy. *PLoS One* 7 (12), e50915.
- Ardeshirpour, Y., Chernomordik, V., Zielinski, R., Capala, J., Griffiths, G., Vaslatiy, O., et al., 2012. *In vivo* fluorescence lifetime imaging monitors binding of specific probes to cancer biomarkers. *PLoS One* 7 (2), e31881.
- Bang, A., Wilhite, T.J., Pike, L.R.G., Cagney, D.N., Aizer, A.A., Taylor, A., et al., 2017. Multicenter Evaluation of the Tolerability of Combined Treatment With PD-1 and CTLA-4 Immune Checkpoint Inhibitors and Palliative Radiation Therapy. *Int. J. Radiat. Oncol. Biol. Phys.* 98 (2), 344–351.
- Bar-Zeev, M., Livney, Y.D., Assaraf, Y.G., 2017. Targeted nanomedicine for cancer therapeutics: Towards precision medicine overcoming drug resistance. *Drug Resist. Updat.* 31, 15–30.
- Barretto, R.P.-J., Ko, T.H., Jung, J.C., Wang, T.J., Capps, G., Waters, A.C., et al., 2011. Time-lapse imaging of disease progression in deep brain areas using fluorescence microendoscopy. *Nat. Med.* 17 (2), 223–228.
- Belcaid, Z., Phallen, J.A., Zeng, J., See, A.P., Mathios, D., Gottschalk, C., et al., 2014. Focal Radiation Therapy Combined with 4-1BB Activation and CTLA-4 Blockade Yields Long-Term Survival and a Protective Antigen-Specific Memory Response in a Murine Glioma Model. *PLoS One* 9 (7), e101764.
- Berghoff, A.S., Kiesel, B., Widhalm, G., Rajky, O., Ricken, G., Wohrer, A., et al., 2015. Programmed death ligand 1 expression and tumor-infiltrating lymphocytes in glioblastoma. *Neuro Oncol.* 17 (8), 1064–1075.
- Berghoff, A.S., Preusser, M., 2015. The inflammatory microenvironment in brain metastases: potential treatment target? *Chin. Clin. Oncol.* 4 (2).
- Berghoff, A.S., Venur, V.A., Preusser, M., Ahluwalia, M.S., 2016. Immune Checkpoint Inhibitors in Brain Metastases: From Biology to Treatment. *Am. Soc. Clin. Oncol. Educ. Book* 35, e116–22.
- Berning, S., Willig, K.I., Steffens, H., Dibaj, P., Hell, S.W., 2012. Nanoscopy in a Living Mouse Brain. *Science* 335 (6068), 551.
- Biswas, Subhra K., 2015. Metabolic Reprogramming of Immune Cells in Cancer Progression. *Immunity* 43 (3), 435–449.
- Boissonnas, A., Fetler, L., Zeelenberg, I.S., Hugues, S., Amigorena, S., 2007. *In vivo* imaging of cytotoxic T cell infiltration and elimination of a solid tumor. *J. Exp. Med.* 204 (2), 345–356.
- Bracci, L., Schiavoni, G., Sistigu, A., Belardelli, F., 2014. Immune-based mechanisms of cytotoxic chemotherapy: implications for the design of novel and rationale-based combined treatments against cancer. *Cell Death Differ.* 21 (1), 15–25.
- Breart, B., Lemaître, F., Celli, S., Bousso, P., 2008. Two-photon imaging of intratumoral CD8+ T cell cytotoxic activity during adoptive T cell therapy in mice. *J. Clin. Invest.* 118 (4), 1390–1397.
- Brown, E.B., Campbell, R.B., Tsuzuki, Y., Xu, L., Carmeliet, P., Fukumura, D., et al., 2001. *In vivo* measurement of gene expression, angiogenesis and physiological function in tumors using multiphoton laser scanning microscopy. *Nat. Med.* 7 (7), 864–868.
- Buchbinder, E.L., Desai, A., 2016. CTLA-4 and PD-1 Pathways: Similarities, Differences, and Implications of Their Inhibition. *Am. J. Clin. Oncol.* 39 (1), 98–106.
- Burke, K., Brown, E., 2014. The Use of Second Harmonic Generation to Image the Extracellular Matrix During Tumor Progression. *Intravital* 3 (3), e984509.
- Butte, M.J., Keir, M.E., Phamduy, T.B., Sharpe, A.H., Freeman, G.J., 2007. Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses. *Immunity* 27 (1), 111–122.
- Caffo, M., Barresi, V., Caruso, G., Cutugno, M., La Fata, G., Venza, M., et al., 2013. Innovative Therapeutic Strategies in the Treatment of Brain Metastases. *Int. J. Mol. Sci.* 14 (1), 2135–2174.
- Campagnola, P.J., Millard, A.C., Terasaki, M., Hoppe, P.E., Malone, C.J., Mohler, W.A., 2002. Three-dimensional high-resolution second-harmonic generation imaging of endogenous structural proteins in biological tissues. *Biophys. J.* 82 (1 Pt 1), 493–508.
- Casey, S.C., Amedei, A., Aquilano, K., Azmi, A.S., Benencia, F., Bhakta, D., et al., 2015. Cancer prevention and therapy through the modulation of the tumor microenvironment. *Semin. Cancer Biol.* 35 (Supplement), S199–S223.
- Chacko, A.M., 2013. Targeted delivery of antibody-based therapeutic and imaging agents to CNS tumors: Crossing the blood-brain-barrier divide. *Expert Opin. Drug Deliv.* 10 (7), 907–926.
- Chmielewski, M., Hombach, A.A., Abken, H., 2014. Of CARs and TRUCKs: chimeric antigen receptor (CAR) T cells engineered with an inducible cytokine to modulate the tumor stroma. *Immunol. Rev.* 257 (1), 83–90.
- Choi, H., Tzeranis, D.S., Cha, J.W., Clémenceau, P., de Jong, S.J., van Geest, L.K., et al., 2012. 3D-resolved fluorescence and phosphorescence lifetime imaging using temporal focusing wide-field two-photon excitation. *Opt. Express* 20 (24), 26219–26235.
- Chtanova, T., Hampton, H.R., Waterhouse, L.A., Wood, K., Tomura, M., Miwa, Y., et al., 2014. Real-time interactive two-photon photoconversion of recirculating lymphocytes for discontinuous cell tracking in live adult mice. *J. Biophotonics* 7 (6), 425–433.
- Cohen, J.V., Kluger, H.M., 2016. Systemic Immunotherapy for the Treatment of Brain Metastases. *Front. Oncol.* 6.
- Coloco, R.J., Martin, P., Kluger, H.M., Yu, J.B., Chiang, V.L., 2016. Does immunotherapy increase the rate of radiation necrosis after radiosurgical treatment of brain metastases? *J. Neurosurg.* 125 (1), 17–23.
- Constantinescu, C.S., Tani, M., Ransohoff, R.M., Wysocka, M., Hilliard, B., Fujioka, T., et al., 2005. Astrocytes as antigen-presenting cells: expression of IL-12/IL-23. *J. Neurochem.* 95 (2), 331–340.
- Curran, M.A., Montalvo, W., Yagita, H., Allison, J.P., 2010. PD-1 and CTLA-4 combination blockade expands infiltrating T cells and reduces regulatory T and myeloid cells within B16 melanoma tumors. *Proc. Natl. Acad. Sci. U. S. A.* 107 (9), 4275–4280.
- De Yébenes, J.G., Gervás, J.J., Iglesias, J., Mena, M.A., Del Río, R.M., Somoza, E., 1982. Biochemical findings in a case of parkinsonism secondary to brain tumor. *Ann. Neurol.* 11 (3), 313–316.
- Degenhardt, K., Mathew, R., Beaudoin, B., Bray, K., Anderson, D., Chen, G., et al., 2006. Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis. *Cancer Cell* 10 (1), 51–64.
- Deguine, J., Breart, B., Lemaître, F., Di Santo, J.P., Bousso, P., 2010. Intravital imaging reveals distinct dynamics for natural killer and CD8(+) T cells during tumor regression. *Immunity* 33 (4), 632–644.
- Dellian, M., Helmlinger, G., Yuan, F., Jain, R.K., 1996. Fluorescence ratio imaging of interstitial pH in solid tumours: effect of glucose on spatial and temporal gradients. *Br. J. Cancer* 74 (8), 1206–1215.
- Demaria, S., Kawashima, N., Yang, A.M., Devitt, M.L., Babb, J.S., Allison, J.P., et al., 2005. Immune-mediated inhibition of metastases after treatment with local radiation and CTLA-4 blockade in a mouse model of breast cancer. *Clin. Cancer Res.* 11 (2 Pt 1), 728–734.

- Deryugina, E.I., Quigley, J.P., 2006. Matrix metalloproteinases and tumor metastasis. *Cancer Metastasis Rev.* 25 (1), 9–34.
- Desai, T.J., Brownfield, D.G., Krasnow, M.A., 2014. Alveolar progenitor and stem cells in lung development, renewal and cancer. *Nature* 507 (7491), 190–194.
- Di Giacomo, A.M., Ascierto, P.A., Pilla, L., Santinami, M., Ferrucci, P.F., Giannarelli, D., et al., 2012. Ipilimumab and fotemustine in patients with advanced melanoma (NIBIT-M1): an open-label, single-arm phase 2 trial. *Lancet Oncol.* 13 (9), 879–886.
- Di Giacomo, A.M., Ascierto, P.A., Queirolo, P., Pilla, L., Ridolfi, R., Santinami, M., et al., 2015. Three-year follow-up of advanced melanoma patients who received ipilimumab plus fotemustine in the Italian Network for Tumor Biotherapy (NIBIT)-M1 phase II study. *Ann. Oncol.* 26 (4), 798–803.
- Di Giacomo, A.M., Margolin, K., 2015. Immune Checkpoint Blockade in Patients With Melanoma Metastatic to the Brain. *Semin. Oncol.* 42 (3), 459–465.
- Diaz-Montero, C.M., Salem, M.L., Nishimura, M.I., Garrett-Mayer, E., Cole, D.J., Montero, A.J., 2009. Increased circulating myeloid-derived suppressor cells correlate with clinical cancer stage, metastatic tumor burden, and doxorubicin-cyclophosphamide chemotherapy. *Cancer Immunol. Immunother.* 58 (1), 49–59.
- Dietzel, S., Pircher, J., Nekolla, A.K., Gull, M., Brandli, A.W., Pohl, U., et al., 2014. Label-free determination of hemodynamic parameters in the microcirculation with third harmonic generation microscopy. *PLoS One* 9 (6), e99615.
- Dye, N.B., Gondi, V., Mehta, M.P., 2015. Strategies for preservation of memory function in patients with brain metastases. *Chin. Clin. Oncol.* 4 (2), 24.
- Egeblad, M., Ewald, A.J., Askautrud, H.A., Truitt, M.L., Welm, B.E., Bainbridge, E., et al., 2008. Visualizing stromal cell dynamics in different tumor microenvironments by spinning disk confocal microscopy. *Dis. Model. Mech.* 1 (2–3), 155–167 discussion 65.
- Ellenbroek, S.I., van Rheejen, J., 2014. Imaging hallmarks of cancer in living mice. *Nat. Rev. Cancer* 14 (6), 406–418.
- Engelhardt, B., Carare, R.O., Bechmann, I., Flügel, A., Laman, J.D., Weller, R.O., 2016. Vascular, glial, and lymphatic immune gateways of the central nervous system. *Acta Neuropathol.* 132, 317–338.
- Evans, C.L., Xie, X.S., 2008. Coherent anti-stokes Raman scattering microscopy: chemical imaging for biology and medicine. *Annu. Rev. Anal. Chem. (Palo Alto Calif)* 1, 883–909.
- Fang, P., Jiang, W., Allen, P., Glitza, I., Guha, N., Hwu, P., et al., 2017. Radiation necrosis with stereotactic radiosurgery combined with CTLA-4 blockade and PD-1 inhibition for treatment of intracranial disease in metastatic melanoma. *J. Neurooncol.*
- Fecci, P.E., Ochiai, H., Mitchell, D.A., Grossi, P.M., Sweeney, A.E., Archer, G.E., et al., 2007. Systemic CTLA-4 blockade ameliorates glioma-induced changes to the CD4+ T cell compartment without affecting regulatory T-cell function. *Clin. Cancer Res.* 13 (7), 2158–2167.
- Fesnak, A.D., June, C.H., Levine, B.L., Engineered, T., 2016. cells: the promise and challenges of cancer immunotherapy. *Nat. Rev. Cancer* 16 (9), 566–581.
- Franceschini, D., Franzese, C., Navarra, P., Ascolese, A.M., De Rose, F., Del Vecchio, M., et al., 2016. Radiotherapy and immunotherapy: Can this combination change the prognosis of patients with melanoma brain metastases? *Cancer Treatment Rev.* 50, 1–8.
- Fridman, W.H., Zitvogel, L., Sautes-Fridman, C., Kroemer, G., 2017. The immune contexture in cancer prognosis and treatment. *Nat. Rev. Clin. Oncol* advance online publication.
- Frieboes, H.B., Smith, B.R., Chuang, Y.L., Ito, K., Roettgers, A.M., Gambhir, S.S., et al., 2013. An integrated computational/experimental model of lymphoma growth. *PLoS Comput. Biol.* 9 (3), e1003008.
- Fukumura, D., Duda, D.G., Munn, L.L., Jain, R.K., 2010. Tumor microvasculature and microenvironment: novel insights through intravital imaging in pre-clinical models. *Microcirculation (New York, NY : 1994)* 17 (3), 206–225.
- Fukumura, D., Jain, R.K., 2007. Tumor microvasculature and microenvironment: targets for anti-angiogenesis and normalization. *Microvasc. Res.* 74 (2–3), 72–84.
- Galon, J., Costes, A., Sanchez-Cabo, F., Kirilovsky, A., Mlecnik, B., Lagorce-Pages, C., et al., 2006. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* 313 (5795), 1960–1964.
- Gershanov, S., Michowiz, S., Toledano, H., Yahav, G., Barinfeld, O., Hirshberg, A., et al., 2017. Fluorescence Lifetime Imaging Microscopy, a Novel Diagnostic Tool for Metastatic Cell Detection in the Cerebrospinal Fluid of Children with Medulloblastoma. *Sci. Rep.* 7 (1), 3648.
- Giedt, R.J., Koch, P.D., Weissleder, R., 2013. Single cell analysis of drug distribution by intravital imaging. *PLoS One* 8 (4), e60988.
- Goldberg, S.B., Gettinger, S.N., Mahajan, A., Chiang, A.C., Herbst, R.S., Sznol, M., et al., 2016. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol.* 17 (7), 976–983.
- Gonen, N., Assaraf, Y.G., 2012. Antifolates in cancer therapy: structure, activity and mechanisms of drug resistance. *Drug Resist. Updat.* 15 (4), 183–210.
- Gordon, S.R., Maute, R.L., Dulken, B.W., Hutter, G., George, B.M., McCracken, M.N., et al., 2017. PD-1 expression by tumour-associated macrophages inhibits phagocytosis and tumour immunity. *Nature* 545 (7655), 495–499.
- Gottesman, M.M., Lavi, O., Hall, M.D., Gillet, J.P., 2016. Toward a Better Understanding of the Complexity of Cancer Drug Resistance. *Annu. Rev. Pharmacol. Toxicol.* 56, 85–102.
- Guan, X., 2015. Cancer metastases: challenges and opportunities. *Acta Pharm. Sin. B* 5 (5), 402–418.
- Hak, S., Reitan, N.K., Haraldseth, O., de Lange Davies, C., 2010. Intravital microscopy in window chambers: a unique tool to study tumor angiogenesis and delivery of nanoparticles. *Angiogenesis* 13 (2), 113–130.
- Hambardzumyan, D., Gutmann, D.H., Kettenmann, H., 2016. The role of microglia and macrophages in glioma maintenance and progression. *Nat. Neurosci.* 19 (1), 20–27.
- Hatiboglu, M.A., Wildrick, D.M., Sawaya, R., 2013. The role of surgical resection in patients with brain metastases. *Ecancermedscience* 7.
- Heller, K.N., Pavlick, A.C., Hodi, F.S., Thompson, J.A., Margolin, K.A., Lawrence, D.P., et al., 2011. Safety and survival analysis of ipilimumab therapy in patients with stable asymptomatic brain metastases. *J. Clin. Oncol.* 29 (Suppl. 15), 8581.
- Helmchen, F., Denk, W., 2005. Deep tissue two-photon microscopy. *Nat. Methods* 2 (12), 932–940.
- Helmlinger, G., Yuan, F., Dellian, M., Jain, R.K., 1997. Interstitial pH and pO₂ gradients in solid tumors in vivo: high-resolution measurements reveal a lack of correlation. *Nat. Med.* 3 (2), 177–182.
- Hervé, F., Ghinea, N., Scherrmann, J.M., 2008. CNS Delivery Via Adsorptive Transcytosis. *AAPS J.* 10 (3), 455–472.
- Herz, J., Filiano, A.J., Smith, A., Yogev, N., Kipnis, J., 2017. Myeloid Cells in the Central Nervous System. *Immunity* 46 (6), 943–956.
- Hoshida, T., Isaka, N., Hagendoorn, J., di Tomaso, E., Chen, Y.L., Pytowski, B., et al., 2006. Imaging steps of lymphatic metastasis reveals that vascular endothelial growth factor-C increases metastasis by increasing delivery of cancer cells to lymph nodes: therapeutic implications. *Cancer Res.* 66 (16), 8065–8075.
- Hottinger, A.F., Pacheco, P., Stupp, R., 2016. Tumor treating fields: a novel treatment modality and its use in brain tumors. *Neuro Oncol.* 18 (10), 1338–1349.
- Iwai, Y., Ishida, M., Tanaka, Y., Okazaki, T., Honjo, T., Minato, N., 2002. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc. Natl. Acad. Sci. U. S. A.* 99 (19), 12293–12297.
- Jain, R.K., 2002. Angiogenesis and lymphangiogenesis in tumors: insights from intravital microscopy. *Cold Spring Harb. Symp. Quant. Biol.* 67, 239–248.
- Jarzyna, P.A., Gianella, A., Skajaa, T., Knudsen, G., Deddens, L.H., Cormode, D.P., et al., 2010. Multifunctional imaging nanoprobes. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* 2 (2), 138–150.
- Jaspers, J.E., Brentjens, R.J., 2017. Development of CAR T cells designed to improve antitumor efficacy and safety. *Pharmacol. Ther.* 178, 83–91.
- Johanns, T., Waqar, S.N., Morgensztern, D., 2016. Immune checkpoint inhibition in patients with brain metastases. *Ann. Transl. Med.* 4 (Suppl. 1).
- Joyce, J.A., Fearon, D.T., 2015. T cell exclusion, immune privilege, and the tumor microenvironment. *Science* 348 (6230), 74–80.
- Kennedy-Darling, Julia, Nolan, Garry P., Goltsev, Yury, Samusik, Nikolay, 2016. Abstract A089: Multiparametric immunofluorescence analysis of the tumor microenvironment using CODEX.
- Kamoun, W.S., Ley, C.D., Farrar, C.T., Duyverman, A.M., Lahdenranta, J., Lacorre, D.A., et al., 2009. Edema control by cediranib, a vascular endothelial growth factor receptor-targeted kinase inhibitor, prolongs survival despite persistent brain tumor growth in mice. *J. Clin. Oncol.* 27 (15), 2542–2552.
- Karagiannis, G.S., Pastoriza, J.M., Wang, Y., Harney, A.S., Entenberg, D., Pignatelli, J., et al., 2017. Neoadjuvant chemotherapy induces breast cancer metastasis through a TMEM-mediated mechanism. *Sci. Transl. Med.* 9 (397).
- Keikhosravi, A., Bredfeldt, J.S., Sagar, A.K., Eliceiri, K.W., 2014. Second-harmonic generation imaging of cancer. *Methods Cell Biol.* 123, 531–546.
- Keir, M.E., Butte, M.J., Freeman, G.J., Sharpe, A.H., 2008. PD-1 and its ligands in tolerance and immunity. *Annu. Rev. Immunol.* 26, 677–704.
- Kessenbrock, K., Plaks, V., Werb, Z., 2010. Matrix metalloproteinases: regulators of the tumor microenvironment. *Cell* 141 (1), 52–67.
- Kienast, Y., von Baumgarten, L., Fuhrmann, M., Klinkert, W.E., Goldbrunner, R., Herms, J., et al., 2010. Real-time imaging reveals the single steps of brain metastasis formation. *Nat. Med.* 16 (1), 116–122.
- Kim, B.M., Eichler, J., Reiser, K.M., Rubenchik, A.M., Da Silva, L.B., 2000. Collagen structure and nonlinear susceptibility: effects of heat, glycation, and enzymatic cleavage on second harmonic signal intensity. *Lasers Surg. Med.* 27 (4), 329–335.
- Knisely, J.P., Yu, J.B., Flanagan, J., Sznol, M., Kluger, H.M., Chiang, V.L., 2012. Radiosurgery for melanoma brain metastases in the ipilimumab era and the possibility of longer survival. *J. Neurosurg.* 117 (2), 227–233.
- Kobat, D., Durst, M.E., Nishimura, N., Wong, A.W., Schaffer, C.B., Xu, C., 2009. Deep tissue multiphoton microscopy using longer wavelength excitation. *Opt. Express* 17 (16), 13354–13364.
- Koffie, R.M., Farrar, C.T., Saidi, L.J., William, C.M., Hyman, B.T., Spires-Jones, T.L., 2011. Nanoparticles enhance brain delivery of blood-brain barrier-impermeable probes for in vivo optical and magnetic resonance imaging. *Proceedings of the National Academy of Sciences of the United States of America* 108 (46), 7–12.
- Koo, T., Kim, I.A., 2016. Radiotherapy and immune checkpoint blockades: a snapshot in 2016. *Radiation oncology journal* 34 (4), 250–259.
- Koyama, S., Akbay, E.A., Li, Y.Y., Herter-Sprie, G.S., Buczkowski, K.A., Richards, W.G., et al., 2016. Adaptive resistance to therapeutic PD-1 blockade is associated with upregulation of alternative immune checkpoints. *Nat. Commun.* 7, 10501.
- Kurokawa, H., Ito, H., Inoue, M., Tabata, K., Sato, Y., Yamagata, K., et al., 2015. High resolution imaging of intracellular oxygen concentration by phosphorescence lifetime. *Sci. Rep.* 5, 10657.
- Kyritsis, A.P., Markoula, S., Levin, V.A., 2012. A systematic approach to the management of patients with brain metastases of known or unknown primary site. *Cancer Chemother. Pharmacol.* 69 (1), 1–13.
- Larkin, J., Chiarion-Sileni, V., Gonzalez, R., Grob, J.J., Cowey, C.L., Lao, C.D., et al., 2015. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N. Eng. J. Med.* 373 (1), 23–34.
- Le, T.T., Huff, T.B., Cheng, J.X., 2009. Coherent anti-Stokes Raman scattering imaging of lipids in cancer metastasis. *BMC Cancer* 9, 42.
- Lee, J., Kotliarova, S., Kotliarov, Y., Li, A., Su, Q., Donin, N.M., et al., 2006. Tumor stem cells derived from glioblastomas cultured in bFGF and EGF more closely mirror the phenotype and genotype of primary tumors than do serum-cultured cell lines. *Cancer*

- Cell 9 (5), 391–403.
- Lee, K.S., Wu, Z., Song, Y., Mitra, S.S., Feroze, A.H., Cheshier, S.H., et al., 2013. Roles of PINK1, mTORC2, and mitochondria in preserving brain tumor-forming stem cells in a noncanonical Notch signaling pathway. *Genes. Dev.* 27 (24), 2642–2647.
- Li, W., Zhang, H., Assaraf, Y.G., Zhao, K., Xu, X., Xie, J., et al., 2016a. Overcoming ABC transporter-mediated multidrug resistance: Molecular mechanisms and novel therapeutic drug strategies. *Drug Resist. Updat.* 27, 14–29.
- Li, Y., Li, F., Jiang, F., Lv, X., Zhang, R., Lu, A., et al., 2016b. A Mini-Review for Cancer Immunotherapy: Molecular Understanding of PD-1/PD-L1 Pathway & Translational Blockade of Immune Checkpoints. *Int. J. Mol. Sci.* 17, 7.
- Lin, J., Er, K.Z.J., Zheng, W., Huang, Z., 2013. Radially polarized tip-enhanced near-field coherent anti-Stokes Raman scattering microscopy for vibrational nano-imaging. *Appl. Phys. Lett.* 103 (8), 083705.
- Lin, X., DeAngelis, L.M., 2015. Treatment of Brain Metastases. *J. Clin. Oncol.* 33 (30), 3475–3484.
- Lorger, M., 2012. Tumor Microenvironment in the Brain. *Cancers (Basel)* 4 (1), 218–243.
- Louveau, A., Smirnov, I., Keyes, T.J., Eccles, J.D., Rouhani, S.J., Peske, J.D., et al., 2015. Structural and functional features of central nervous system lymphatic vessels. *Nature* 523 (7560), 337–341.
- Lu, W., Su, J., Kim, L.S., Bucana, C.D., Donawho, C., He, J., et al., 2003. Active specific immunotherapy against occult brain metastasis. *Cancer Res.* 63 (6), 1345–1350.
- Margolin, K., Ernstoff, M.S., Hamid, O., Lawrence, D., McDermott, D., Puzanov, I., et al., 2012. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol.* 13 (5), 459–465.
- Massague, J., Obenauf, A.C., 2016. Metastatic colonization by circulating tumour cells. *Nature* 529 (7586), 298–306.
- Maurin, M., Stephan, O., Vial, J.C., Marder, S.R., van der Sanden, B., 2011. Deep in vivo two-photon imaging of blood vessels with a new dye encapsulated in pluronic nanomicelles. *J. Biomed. Opt.* 16 (3), 036001.
- Melder, R.J., Salehi, H.A., Jain, R.K., 1995. Interaction of activated natural killer cells with normal and tumor vessels in cranial windows in mice. *Microvasc. Res.* 50 (1), 35–44.
- Mikucki, M.E., Fisher, D.T., Matsuzaki, J., Skitzki, J.J., Gaulin, N.B., Muhitch, J.B., et al., 2015. Non-redundant requirement for CXCR3 signalling during tumoricidal T-cell trafficking across tumour vascular checkpoints. *Nat. Commun.* 6, 7458.
- Mittal, S., Klinger, N.V., Michelhaugh, S.K., Barger, G.R., Pannullo, S.C., Juhász, C., 2017. Alternating electric tumor treating fields for treatment of glioblastoma: rationale, preclinical, and clinical studies. *J. Neurosurg.* 0 (0), 1–8.
- Moalli, F., Proulx, S.T., Schwendener, R., Detmar, M., Schlapbach, C., Stein, J.V., 2015. Intravital and whole-organ imaging reveals capture of melanoma-derived antigen by lymph node subcapsular macrophages leading to widespread deposition on follicular dendritic cells. *Front. Immunol.* 6, 114.
- Monsky, W.L., Mouta Carreira, C., Tsuzuki, Y., Gohongi, T., Fukumura, D., Jain, R.K., 2002. Role of host microenvironment in angiogenesis and microvascular functions in human breast cancer xenografts: mammary fat pad versus cranial tumors. *Clin. Cancer Res.* 8 (4), 1008–1013.
- Muradashvili, N., Tyagi, R., Lominadze, D., 2012. A Dual-Tracer Method for Differentiating Transendothelial Transport from Paracellular Leakage in Vivo and in Vitro. *Front. Physiol.* 3.
- Murdoch, C., Muthana, M., Coffelt, S.B., Lewis, C.E., 2008. The role of myeloid cells in the promotion of tumour angiogenesis. *Nat. Rev. Cancer* 8 (8), 618–631.
- Naidoo, J., Page, D.B., Wolchok, J.D., 2014. Immune modulation for cancer therapy. *Br. J. Cancer* 111 (12), 2214–2219.
- NCT02085070. MK-3475 in Melanoma and NSCLC Patients With Brain Metastases. < <https://clinicaltrials.gov/ct2/show/NCT02085070?term=NCT02085070&rank=1> > .
- Newick, K., O'Brien, S., Moon, E., Albelda, S.M., 2017. CAR T. Cell Therapy for Solid Tumors. *Annu. Rev. Med.* 68, 139–152.
- Nobis, M., Carragher, N.O., McGhee, E.J., Morton, J.P., Sansom, O.J., Anderson, K.I., et al., 2013. Advanced intravital subcellular imaging reveals vital three-dimensional signalling events driving cancer cell behaviour and drug responses in live tissue. *FEBS J.* 280 (21), 5177–5197.
- Oheim, M., Michael, D.J., Geisbauer, M., Madsen, D., Chow, R.H., 2006. Principles of two-photon excitation fluorescence microscopy and other nonlinear imaging approaches. *Adv. Drug Deliv. Rev.* 58 (7), 788–808.
- Ohkubo, C., Bigos, D., Jain, R.K., 1991. Interleukin 2 induced leukocyte adhesion to the normal and tumor microvascular endothelium in vivo and its inhibition by dextran sulfate: implications for vascular leak syndrome. *Cancer Res.* 51 (5), 1561–1563.
- Oron, D., Yelin, D., Tal, E., Raz, S., Fachima, R., Silberberg, Y., 2004. Depth-resolved structural imaging by third-harmonic generation microscopy. *J. Struct. Biol.* 147 (1), 3–11.
- Osswald, M., Winkler, F., 2013. Insights into cell-to-cell and cell-to-blood-vessel communications in the brain: in vivo multiphoton microscopy. *Cell Tissue Res.* 352 (1), 149–159.
- Ott, P.A., Hodi, F.S., Kaufman, H.L., Wigginton, J.M., Wolchok, J.D., 2017. Combination immunotherapy: a road map. *J. Immunother.* Cancer 5 (1), 16.
- Owen, S., Souhami, L., 2014. The Management of Brain Metastases in Non-Small Cell Lung Cancer. *Front. Oncol.* 4.
- Pattabiraman, D.R., Weinberg, R.A., 2014. Tackling the cancer stem cells – what challenges do they pose? *Nat. Rev. Drug Discov.* 13 (7), 497–512.
- Peinado, H., Zhang, H., Matei, I.R., Costa-Silva, B., Hoshino, A., Rodrigues, G., et al., 2017. Pre-metastatic niches: organ-specific homes for metastases. *Nat. Rev. Cancer* 17 (5), 302–317.
- Pilon-Thomas, S., Mackay, A., Vohra, N., Mule, J.J., 2010. Blockade of programmed death ligand 1 enhances the therapeutic efficacy of combination immunotherapy against melanoma. *J. Immunol. (Baltimore, Md. : 1950)* 184 (7), 3442–3449.
- Pitt, J.M., Vétizou, M., Daillère, R., Roberti, M.P., Yamazaki, T., Routy, B., et al., 2017. Resistance Mechanisms to Immune-Checkpoint Blockade in Cancer: Tumor-Intrinsic and — Extrinsic Factors. *Immunity* 44 (6), 1255–1269.
- Pittet, M.J., Weissleder, R., 2011. Intravital Imaging. *Cell* 147 (5), 983–991.
- Pittet, M.J., Weissleder, R., 2016. Intravital Imaging. *Cell* 147 (5), 983–991.
- Pons-Salort, M., van der Sanden, B., Juhem, A., Popov, A., Stéphanou, A., 2012. A Computational Framework to Assess the Efficacy of Cytotoxic Molecules and Vascular Disrupting Agents against Solid Tumours. *Math Model Nat. Phenom.* 7 (1), 49–77.
- Pope, I., Payne, L., Zorinians, G., Thomas, E., Williams, O., Watson, P., et al., 2014. Coherent anti-Stokes Raman scattering microscopy of single nanodiamonds. *Nat Nano* 9 (11), 940–946.
- Prins, R.M., Vo, D.D., Khan-Farooqi, H., Yang, M.Y., Soto, H., Economou, J.S., et al., 2006. NK and CD4 cells collaborate to protect against melanoma tumor formation in the brain. *J. Immunol. (Baltimore Md. : 1950)* 177 (12), 8448–8455.
- Provenzano, P.P., Inman, D.R., Eliceiri, K.W., Knittel, J.G., Yan, L., Rueden, C.T., et al., 2008. Collagen density promotes mammary tumor initiation and progression. *BMC Med.* 6, 11.
- Puhalla, S., Elmquist, W., Freyer, D., Kleinberg, L., Adkins, C., Lockman, P., et al., 2015. Unsantifying the sanctuary: challenges and opportunities with brain metastases. *Neuro Oncol.* 17 (5), 639–651.
- Pukrop, T., Dehghani, F., Chuang, H.N., Lohaus, R., Bayanga, K., Heermann, S., et al., 2010. Microglia promote colonization of brain tissue by breast cancer cells in a Wnt-dependent way. *Glia* 58 (12), 1477–1489.
- Pyne, S., Hu, X., Wang, K., Rossin, E., Lin, T.I., Maier, L.M., et al., 2009. Automated high-dimensional flow cytometric data analysis. *Proc. Natl. Acad. Sci. U. S. A.* 106 (21), 8519–8524.
- Qiu, P., Simonds, E.F., Bendall, S.C., Gibbs Jr., K.D., Bruggner, R.V., Linderman, M.D., et al., 2011. Extracting a cellular hierarchy from high-dimensional cytometry data with SPADE. *Nat. Biotechnol.* 29 (10), 886–891.
- Quail, D.F., Joyce, J.A., 2017. The Microenvironmental Landscape of Brain Tumors. *Cancer Cell* 31 (3), 326–341.
- Reardon, D.A., Gokhale, P.C., Klein, S.R., Ligon, K.L., Rodig, S.J., Ramkissoon, S.H., et al., 2016. Glioblastoma Eradication Following Immune Checkpoint Blockade in an Orthotopic, Immunocompetent Model. *Cancer Immunol. Res.* 4 (2), 124–135.
- Regina, A., Demeule, M., Laplante, A., Jodoin, J., Dagenais, C., Berthelet, F., et al., 2001. Multidrug resistance in brain tumors: roles of the blood-brain barrier. *Cancer Metastasis Rev.* 20 (1–2), 13–25.
- Reits, E.A., Hodge, J.W., Herbets, C.A., Groothuis, T.A., Chakraborty, M., et al., 2006. Radiation modulates the peptide repertoire, enhances MHC class I expression, and induces successful antitumor immunotherapy. *J. Exp. Med.* 203 (5), 1259–1271.
- Ricard, C., Debarbieux, F.C., 2014. Six-color intravital two-photon imaging of brain tumors and their dynamic microenvironment. *Front. Cell. Neurosci.* 8, 57.
- Riella, L.V., Paterson, A.M., Sharpe, A.H., Chandraker, A., 2012. Role of the PD-1 Pathway in the Immune Response. *Am. J. Transplant.* 12 (10), 2575–2587.
- Rodriguez, P.C., Ochoa, A.C., Al-Khami, A.A., 2017. Arginine Metabolism in Myeloid Cells Shapes Innate and Adaptive Immunity. *Front. Immunol.* 8, 93.
- Ryu, J.H., Koo, H., Sun, I.C., Yuk, S.H., Choi, K., Kim, K., et al., 2012. Tumor-targeting multi-functional nanoparticles for theragnosis: new paradigm for cancer therapy. *Adv. Drug Deliv. Rev.* 64 (13), 1447–1458.
- Shevach, E.M., 2011. Biological functions of regulatory T cells. *Adv. Immunol.* 112, 137–176.
- Sica, A., Bronte, V., 2007. Altered macrophage differentiation and immune dysfunction in tumor development. *J. Clin. Invest.* 117 (5), 1155–1166.
- Silk, A.W., Bassetti, M.F., West, B.T., Tsien, C.I., Lao, C.D., 2013. Ipilimumab and radiation therapy for melanoma brain metastases. *Cancer Med.* 2 (6), 899–906.
- Sindoni, A., Minutoli, F., Ascenti, G., Pergolizzi, S., 2017. Combination of immune checkpoint inhibitors and radiotherapy: Review of the literature. *Crit. Rev. Oncol. Hematol.* 113, 63–70.
- Smith, B.R., Gambhir, S.S., 2017. Nanomaterials for In Vivo Imaging. *Chem. Rev.* 117 (3), 901–986.
- Smith, B.R., Ghosn, E.E., Rallapalli, H.J.A., P, Larson, T., Herzenberg, L.A., et al., 2014. Selective uptake of single-walled carbon nanotubes by circulating monocytes for enhanced tumour delivery. *Nat. Nanotechnol.* 9 (6), 481–487.
- Smith, B.R., Kempen, P., Bouley, D., Xu, A., Liu, Z., Melosh, N., et al., 2012. Shape Matters: Intravital Microscopy Reveals Surprising Geometrical Dependence for Nanoparticles in Tumor Models of Extravasation. *Nano Lett.* 12 (7), 3369–3377.
- Smith, B.R., Zavaleta, C., Rosenberg, J., Tong, R., Ramunas, J., Liu, Z., et al., 2013. High-resolution, serial intravital microscopic imaging of nanoparticle delivery and targeting in a small animal tumor model. *Nano Today* 8 (2).
- Staudt, M., Lasithiotakis, K., Leiter, U., Meier, F., Eigentler, T., Bamberg, M., et al., 2010. Determinants of survival in patients with brain metastases from cutaneous melanoma. *Br. J. Cancer* 102 (8), 1213–1218.
- Stoletov, K., Strnad, J., Zardouian, E., Momiyama, M., Park, F.D., Kelber, J.A., et al., 2013. Role of connexins in metastatic breast cancer and melanoma brain colonization. *J. Cell Sci.* 126 (4), 904–913.
- Sukumar, M., Kishton, R.J., Restifo, N.P., 2017. Metabolic reprogramming of anti-tumor immunity. *Curr. Opin. Immunol.* 46, 14–22.
- Sun, Y., Phipps, J., Elson, D.S., Stoy, H., Tinsling, S., Meier, J., et al., 2009. Fluorescence lifetime imaging microscopy: in vivo application to diagnosis of oral carcinoma. *Opt. Lett.* 34 (13), 2081–2083.
- Tabatabaei, P., Visse, E., Bergström, P., Brännström, T., Siesjö, P., Bergenheim, A.T., 2017. Radiotherapy induces an immediate inflammatory reaction in malignant glioma: a clinical microdialysis study. *J. Neurooncol.* 131 (1), 83–92.
- Tan, A.C., Heimberger, A.B., Menzies, A.M., Pavlakis, N., Khasraw, M., 2017. Immune Checkpoint Inhibitors for Brain Metastases. *Curr. Oncol. Rep.* 19 (6), 38.
- Tazi, K., Hathaway, A., Chiuzan, C., Shirai, K., 2015. Survival of melanoma patients with

- brain metastases treated with ipilimumab and stereotactic radiosurgery. *Cancer Med.* 4 (1), 1–6.
- Theer, P., Hasan, M.T., Denk, W., 2003. Two-photon imaging to a depth of 1000 microm in living brains by use of a Ti:Al₂O₃ regenerative amplifier. *Opt. Lett.* 28 (12), 1022–1024.
- Torcellan, T., Stolp, J., Chtanova, T., 2017. In Vivo Imaging Sheds Light on Immune Cell Migration and Function in Cancer. *Front. Immunol.* 8, 309.
- Tserevelakis, G.J., Megalou, E.V., Filippidis, G., Petanidou, B., Fotakis, C., Tavernarakis, N., 2014. Label-free imaging of lipid depositions in *C. elegans* using third-harmonic generation microscopy. *PLoS One* 9 (1), e84431.
- Twyman-Saint Victor, C., Rech, A.J., Maity, A., Rengan, R., Pauken, K.E., Stelekati, E., et al., 2015. Radiation and dual checkpoint blockade activate non-redundant immune mechanisms in cancer. *Nature* 520 (7547), 373–377.
- Uckermann, O., Galli, R., Tamosaityte, S., Leipnitz, E., Geiger, K.D., Schackert, G., et al., 2014. Label-free delineation of brain tumors by coherent anti-Stokes Raman scattering microscopy in an orthotopic mouse model and human glioblastoma. *PLoS One* 9 (9), e107115.
- Vakoc, B.J., Lanning, R.M., Tyrrell, J.A., Padera, T.P., Bartlett, L.A., Stylianopoulos, T., et al., 2009. Three-dimensional microscopy of the tumor microenvironment in vivo using optical frequency domain imaging. *Nat. Med.* 15 (10), 1219–1223.
- Vakoc, B.J., Yun, S.H., de Boer, J.F., Tearney, G.J., Bouma, B.E., 2005. Phase-resolved optical frequency domain imaging. *Opt. Express* 13 (14), 5483–5493.
- van Munster, E.B., Gadella, T.W.J., 2005. Fluorescence Lifetime Imaging Microscopy (FLIM). *Adv. Biochem. Eng. Biotechnol.* 95, 143–175.
- Vasilevko, V., Ghochikyan, A., Holterman, M.J., Agadjanyan, M.G., 2002. CD80 (B7-1) and CD86 (B7-2) are functionally equivalent in the initiation and maintenance of CD4+ T-cell proliferation after activation with suboptimal doses of PHA. *DNA Cell Biol.* 21 (3), 137–149.
- Verma, N., Cowperthwaite, M.C., Burnett, M.G., Markey, M.K., 2013. Differentiating tumor recurrence from treatment necrosis: a review of neuro-oncologic imaging strategies. *Neuro Oncol.* 15 (5), 515–534.
- Vishwanath, K., Yuan, H., Barry, W.T., Dewhirst, M.W., Ramanujam, N., 2009. Using Optical Spectroscopy to Longitudinally Monitor Physiological Changes within Solid Tumors. *Neoplasia (New York, NY)* 11 (9), 889–900.
- Walrath, J.C., Hawes, J.J., Van Dyke, T., Reilly, K.M., 2010. Genetically Engineered Mouse Models in Cancer Research. *Adv. Cancer Res.* 106, 113–164.
- Weber, J., Thompson, J.A., Hamid, O., Minor, D., Amin, A., Ron, I., et al., 2009. A randomized, double-blind, placebo-controlled, phase II study comparing the tolerability and efficacy of ipilimumab administered with or without prophylactic budesonide in patients with unresectable stage III or IV melanoma. *Clin. Cancer Res.* 15 (17), 5591–5598.
- Weber, J.S., Amin, A., Minor, D., Siegel, J., Berman, D., O'Day, S.J., 2011. Safety and clinical activity of ipilimumab in melanoma patients with brain metastases: retrospective analysis of data from a phase 2 trial. *Melanoma Res.* 21 (6), 530–534.
- Weichselbaum, R.R., Liang, H., Deng, L., Fu, Y.X., 2017. Radiotherapy and immunotherapy: a beneficial liaison? *Nat. Rev. Clin. Oncol.* 14 (6), 365–379.
- Weigelin, B., Bakker, G.J., Friedl, P., 2016. Third harmonic generation microscopy of cells and tissue organization. *J. Cell Sci.* 129 (2), 245–255.
- Whiteside, T.L., 2008. The tumor microenvironment and its role in promoting tumor growth. *Oncogene* 27 (45), 5904–5912.
- Wijdeven, R.H., Pang, B., Assaraf, Y.G., Neefjes, J., 2016. Old drugs, novel ways out: Drug resistance toward cytotoxic chemotherapeutics. *Drug Resist. Updat.* 28, 65–81.
- Wilson, E.H., Weninger, W., Hunter, C.A., 2010. Trafficking of immune cells in the central nervous system. *J. Clin. Invest.* 120 (5), 1368–1379.
- Witte, S., Negrean, A., Lodder, J.C., de Kock, C.P., Testa Silva, G., Mansvelter, H.D., et al., 2011. Label-free live brain imaging and targeted patching with third-harmonic generation microscopy. *Proc. Natl. Acad. Sci. U. S. A.* 108 (15), 5970–5975.
- Wu, N.Z., Klitzman, B., Dodge, R., Dewhirst, M.W., 1992. Diminished leukocyte-endothelium interaction in tumor microvessels. *Cancer Res.* 52 (15), 4265–4268.
- Wu, Q., Allouch, A., Martins, I., Brenner, C., Modjtahedi, N., Deutsch, E., et al., 2017. Modulating Both Tumor Cell Death and Innate Immunity Is Essential for Improving Radiation Therapy Effectiveness. *Front. Immunol.* 8.
- Wyckoff, J.B., Wang, Y., Lin, E.Y., Li, J.F., Goswami, S., Stanley, E.R., et al., 2007. Direct visualization of macrophage-assisted tumor cell intravasation in mammary tumors. *Cancer Res.* 67 (6), 2649–2656.
- Yao, Y., Liu, R., Shin, M.S., Trentalange, M., Allore, H., Nassar, A., et al., 2014. CyTOF supports efficient detection of immune cell subsets from small samples. *J. Immunol. Methods* 415, 1–5.
- Yoshimoto, Y., Suzuki, Y., Mimura, K., Ando, K., Oike, T., Sato, H., et al., 2014. Radiotherapy-induced anti-tumor immunity contributes to the therapeutic efficacy of irradiation and can be augmented by CTLA-4 blockade in a mouse model. *PLoS One* 9 (3), e92572.
- Yu, S., Li, A., Liu, Q., Li, T., Yuan, X., Han, X., et al., 2017. Chimeric antigen receptor T cells: a novel therapy for solid tumors. *J. Hematol. Oncol.* 10 (1), 78.
- Yun, S.H., Tearney, G.J., de Boer, J.F., Iftimia, N., Bouma, B.E., 2003. High-speed optical frequency-domain imaging. *Opt. Express* 11 (22), 2953–2963.
- Zeng, J., See, A.P., Phallen, J., Jackson, C.M., Belcaid, Z., Ruzzevick, J., et al., 2013. Anti-PD-1 blockade and stereotactic radiation produce long-term survival in mice with intracranial gliomas. *Int. J. Radiat. Oncol. Biol. Phys.* 86 (2), 343–349.
- Zhitomirsky, B., Assaraf, Y.G., 2016. Lysosomes as mediators of drug resistance in cancer. *Drug Resistance Updat.* 24, 23–33.
- Zhou, Z.N., Boimel, P.J., Segall, J.E., 2011. Tumor-stroma In vivo assays and intravital imaging to study cell migration and metastasis. *Drug Discov. Today Dis. Models* 8 (2–3), 95–112.
- Zipfel, W.R., Williams, R.M., Webb, W.W., 2003. Nonlinear magic: multiphoton microscopy in the biosciences. *Nat. Biotechnol.* 21 (11), 1369–1377.
- Zitvogel, L., Apetoh, L., Ghiringhelli, F., Kroemer, G., 2008. Immunological aspects of cancer chemotherapy. *Nat. Rev. Immunol.* 8 (1), 59–73.